

Italian Guidelines for the diagnosis and treatment of Fetal Alcohol Spectrum Disorders: cognitive and behavioral deficits

GIOVANNA CORIALE¹, ARIANNA BARZACCHI², MAURO CECCANTI², LUIGI TARANI³, SIMONA GENCARELLI¹, MARIA GIUSEPPA ELMO⁴, MARTINO MISTRETTA¹, MARISA PATRIZIA MESSINA³, SIMONE DE PERSIS⁵, MARCO FIORE⁶, DANIELA FIORENTINO⁷; INTERDISCIPLINARY STUDY GROUPS* SAPIENZA, AIDEFAD, SITAC, SIFASD, FIMMG-LAZIO, SIPPS, SIMPESV, CIPE

¹CRARL Lazio, ASL Roma 1, Italy; ²SITAC - Società Italiana per il Trattamento dell'Alcolismo e le sue Complicanze, Rome, Italy; ³Department of Maternal Infantile and Urological Sciences, Sapienza University of Rome, Italy; ⁴Department of Mental Health, ASL Roma 1, Italy; ⁵Department of Mental Health, Rieti, Italy; ⁶Institute of Biochemistry and Cell Biology (IBBC-CNR), Department of Sensory Organs, Sapienza University of Rome, Italy; ⁷ASL Rieti, Italy.

Summary. Objective. Fetal alcohol spectrum disorders (FASDs) refer to a group of clinical conditions that occur in a person exposed to alcohol before birth. Neuroimaging shows abnormalities in brain structure, cortical development, white matter microstructure, and functional connectivity in individuals with FASD. These abnormalities modify the normal developmental trajectories resulting in deficits in cognition and behavior across several domains, including general intelligence, memory, language, attention, learning, visuospatial abilities, executive functioning, fine and gross motor skills, and social and adaptive functioning. This paper provides a review of the cognitive and behavioral outcomes of prenatal alcohol exposure. Updates data on FASD-specific neurobehavioral profile and its potential as a diagnostic tool will then be presented.

Key words. ARND, behavior problems, cognitive development, FASD, neurobehavioral profile.

Introduction

Fetal alcohol spectrum disorders (FASD) refer to a broad spectrum of physical, mental, behavioral, and cognitive abnormalities occurring in individuals with a history of prenatal alcohol exposure (PAE) by the mother or the father¹⁻³ including fetal alcohol syndrome (FAS), partial FAS, alcohol-related neurodevelopmental disorder (ARND) and alcohol-related birth defects (ARBDs)⁴. FASD is associated with several neurobehavioral impairments including general intelligence, adaptive function, attention and executive function, motor function, social cognition, visual-spatial functioning, verbal and nonverbal learning as well as internalizing and externalizing behaviors⁵. Morton and Frith's model can be used to describe

Linee guida italiane per la diagnosi e il trattamento dei disturbi dello spettro feto-alcolico: deficit cognitivi e comportamentali.

Sommario. Obiettivo I disturbi dello spettro feto-alcolico (FASD) fanno riferimento a un gruppo di condizioni cliniche causate dall'esposizione all'alcol durante la gravidanza. Gli studi di neuroimmagini stanno fornendo evidenze sulla presenza di anomalie nella struttura cerebrale, nello sviluppo corticale, nella microstruttura della materia bianca e nella connettività funzionale nei soggetti con FASD. Queste anomalie modificano le normali traiettorie di sviluppo, con conseguenti deficit che influenzano molteplici aspetti della cognizione e del comportamento in diversi domini, tra cui l'intelligenza generale, la memoria, il linguaggio, l'attenzione, l'apprendimento, le abilità visuo-spaziali, il funzionamento esecutivo, le abilità motorie fini e grossolane e il funzionamento sociale e adattivo. Questo documento fornisce una revisione degli esiti cognitivi e comportamentali dell'esposizione prenatale all'alcol. Verranno poi presentati i dati aggiornati sul profilo neurocomportamentale specifico della FASD e il suo potenziale come strumento diagnostico.

Parole chiave. ARND, FASD, problemi comportamentali, profilo neurocomportamentale, sviluppo cognitivo.

FASD-related neurocognitive and behavioral disorders according to three levels of analysis: neurobiological, cognitive, and behavioral⁶.

Neurobiology: structural and functional alterations

A baby with FASD may be born with a head and brain significantly smaller than a normal-sized baby of the same gender and age. Concerning FASD, microcephaly reflects structural damage to the brain. Studies of brain structure using neuroimaging techniques have shown the global and focal effects that prenatal alcohol exposure has on the brain demonstrating reductions in the volume of various brain structures, including total brain, corpus callosum, cerebellum,

and basal ganglia^{7,8}. Alterations in the corpus callosum, often damaged by PAE, have been associated with deficits in several domains of neuropsychological function such as motor function⁹, attention¹⁰, verbal learning¹¹, and executive function¹². Notably, the caudate was the first reported region to have a lower volume associated with prenatal alcohol exposure¹³, a finding that has been consistently reported in the literature¹⁴⁻¹⁶. In addition, the volume of the parietal, temporal, and frontal lobes have all shown locally decreased volume in FASD children and adolescents relative to controls^{14,17}. Very interesting the results of several studies that used task-based fMRI designs have examined the brain function of subjects while performing a cognitive task. Altered brain function has been observed in FASD during various cognitive tasks, including response inhibition, mathematics, and number processing, working memory and verbal learning¹⁸ showing more regional functional demand in children/adolescents with FASD.

According to well-documented structural and microstructural abnormalities in PAE, recent studies suggest that also functional connectivity (FC) may be altered in individuals with FASD¹⁹. In FASD, alterations have been found both within and between most cognition-related networks^{20,21}.

Cognitive impairments

Cognitive deficits affect different domains of cognition^{5,22-27}. The domains are closely related to each other so the alteration of one affects all the others and are hierarchical, with the bottom referring to more basic sensory and perceptual processes and the top referring to elements of executive functioning and cognitive control²⁸. The executive tasks often involve coordinating multiple sensory, perceptual, attentional, and other less complex functions, while simple sensory tasks demand minimal higher-level processing. The top-down vs bottom-up perspective will be used to describe cognitive impairment in FASD.

SENSATION AND PERCEPTION

The sensory information is processed and integrated into the domain of perception. One of the important goals of perception is the identification of previously experienced objects from sensory information and can be assessed in terms of the ability to recognize objects, sounds, and perceptual field in its entirety. It has been found that children with FASDs have a high prevalence of ophthalmologic abnormalities, as subnormal visual acuity (VA), optic nerve hypoplasia (ONH), retinal vascular malformations; refractive errors, and strabismus are well documented in FASDs^{29,30}.

Thus, an ophthalmological assessment should be an integral part of the FASD diagnostic workup, to better understand the nature of sensory and perceptual problems if they were present³¹. Sensory processing (also called sensory integration) refers to the way the individual's nervous system receives signals from sensory modalities and transforms them into motor and behavioral responses while the ability to identify a meaningful stimulus falls under the domain of perception³². Sensory Processing Disorder (SPD) leads to inappropriate behavioral reactions to environmental stimuli making it difficult to organize sensory signals into appropriate responses³³ and since it is commonly comorbid with FASDs, early recognition, diagnosis, and referral for SPDs is important when diagnosing FASD. The occurrence of visual perception problems (VPPs) in children with FASD has previously been reported and their persistence into adulthood has not yet been investigated²⁹. Deficits related to FASD have been described in visual perception³⁴, visual construction³⁵, and feature processing³⁶ (figure 1).

MOTOR SKILLS AND VISUOSPATIAL CONSTRUCTION

Motor skills refer to fine motor abilities, including manual dexterity and motor speed, as well as reaction time, and gross motor skills such as balance. The prevalence of serious fine and gross motor deficits among children diagnosed with FASD is estimated at 51% and 37%, respectively, with clinically important developmental delays more in fine motor skills than in gross motor skills³⁷. Severe fine motor impairment can adversely affect the child's ability to meet daily activity demands, making those difficult to operate and not supporting independence in self-care tasks such as dressing, eating, brushing hair, and cleaning teeth; academic skills including handwriting, drawing, and using scissors; and participation in play and social activities³⁷. Visual-motor integration (VMI) can be defined as the extent to which visual perception and finger-hand movements are well coordinated. Testing in this domain examines the accuracy of an individual's perception of stimuli and their related ability to appropriately manipulate objects in the environment³⁸. Studies have found that VMI skills have a positive influence on fine motor development, handwriting performance, and academic achievements³⁹. Children with FASD have been found poor in VMI and fine motor coordination skills⁴⁰⁻⁴⁴.

The capacity to see an object or image as a composite of parts and then to construct the original from these parts is referred to as visuospatial constructive cognition⁴⁵. Children with FASD seem to have difficulty seeing the complex object as a whole or 'gestalt,' and integrating pieces of the design into a cohesive whole⁴⁶. Deficits include problems with basic figure copying, spatial learning, spatial working memory,




Figure 1. The three primary sensory processing disorders descriptions.

spatial recall, visual-spatial reasoning, visual-perceptual matching (e.g., matching complex geometric shapes), and sustained visual attention^{23,47,48}. Since early motor skill, VMI and visuospatial deficits have been found to have an important and negative impact on functional abilities and academic achievement, physiotherapists and occupational therapists should support the diagnosis of FASD by conducting assessments of the motor domain and providing strategies^{49,50}.

MEMORY AND LEARNING

Memory is an important cognitive function that allows us to acquire, retain, and recover data. Three main processes that characterize how memory works are encoding, storage, and retrieval (or recall). Memory isn't a single unitary phenomenon and is composed of several distinct but interrelated constituent processes and systems⁵¹. There are three major types of human memory: working memory, declarative memory (explicit), and non-declarative memory (implicit). Working memory is a limited capacity store for retaining information for a brief period while performing mental operations on that information. It is critical for a variety of activities at school, from complex subjects such as reading comprehension to mental arithmetic⁵¹. Clinical studies have reported learning and memory deficits in children with

heavy PAE. These impairments range across specific aspects of learning and memory, including verbal and nonverbal skills⁵²⁻⁵⁴. Surprisingly, memory impairment in individuals with FASD has been found the result of erroneous encoding processes rather than retrieval impairments^{5,55-57}. Researchers have obtained evidence of working memory deficits in children with FASD^{26,58}. Impairments in spatial working memory are also evident, with deficits becoming more significant as the task increases in complexity⁵⁹.

LANGUAGE

Language skills include receptive (comprehension) and productive (speech) abilities as well as the ability to access semantic memory, to identify objects with a name, and to respond to verbal instructions with behavioral acts²⁸. Language skills are assessed with measures of fluency by asking to name as many animals as possible, object naming and responding to instructions. Evidence from the longitudinal studies suggests that PAE causes delays in receptive and expressive communication⁶⁰. Various language impairments have been described including problems in speech discrimination, comprehension, syntax development, and prosodic features. However, a child with FASD may present adequate superficial speech skills but also an impairment in pragmatic language use, resulting in poor peer relationships and thus sig-

nificant social problems⁶¹. Preschool children with fetal alcohol exposure have been reported as having social communication deficits and adverse social interactive experiences¹⁴.

ATTENTION

Over the years, researchers have identified various types of attention in psychology. Selective attention is the process of attending to information that is relevant and important and ignoring other non-relevant information. Selective attention tasks often provide distracting information and request the examinee to attend specifically to the relevant information. Instead, sustained attention is the ability to sustain attention over time. Tasks measuring sustained attention as continuous performance tasks often require the detection of simple stimuli presented infrequently during a stream of other stimuli⁶².

When the focus is on two or more things at the same time, divided attention is used. This ability is also known as multitasking and is a crucial factor in the academic setting. The Stroop test is one of the most important tests used to measure divided attention^{63,64}. These studies support the idea that there is a behavioral impairment in the ability to maintain focused alertness in perceiving a signal and in the ability to allocate attentional resources, or the ability to shift attention from one task to another when appropriate^{65,66}. Additionally, attending to visual information appears to be more severely impaired compared to auditory information^{10,67}. Coles et al.⁶⁸ found a greater difficulty in Encode (i.e., the capacity to hold information temporarily in memory while performing a mental operation upon it) and Shift (e.g., the ability to shift attention from one stimulus dimension to another in a flexible manner) components in FASD group. In contrast, the ADHD group displayed greater difficulty on tests assessing Focus (i.e., the capacity to concentrate on a particular task) and Sustain (i.e., the ability to stay on a task) components.

EXECUTIVE FUNCTIONING AND INFORMATION PROCESSING

Executive functions (EFs) are a set of cognitive skills that are needed for self-control and managing behaviors. These skills include high-order cognitive abilities such as working memory, inhibitory control, cognitive flexibility, planning, reasoning, and problem-solving allowing people to follow directions, focus, control emotions, and attain goals⁶⁹. The executive system manages and controls other cognitive abilities (for example attention and memory) and allows individuals to change their overlearned behavioral patterns when they become unsatisfactory⁷⁰. This makes also possible to adapt to new and complex daily

life situations⁶⁶. Consequently, deficits in EFs have important impacts on everyday life, which include poor goal-directed behavior and impaired social functioning⁷¹. Traditionally those have been associated with frontal lobe functioning, but posterior and subcortical regions also play a crucial role in EF processing, especially in the integration of sensory information and emotion. Frontal circuits are particularly vulnerable to PAE as shown by fMRI studies examining executive functioning⁷². The disruption in executive functioning is one of the most important distinctive signs of FASD, characterized by problems with self-control and self-regulation^{59,73-75}. Weaknesses are most consistent for measures of planning, fluency, and set-shifting⁷⁶. EF deficits have been found evident before the age of 6 years in children with FASD, present across all the spectrum and impaired in children with more severe forms of FASD and/or lower IQs and are related to difficulties in daily functioning in children with FASD, limiting independence (routine activities like getting dressed require sequencing skills) and disrupting social interactions^{77,78}. Executive Functioning has been categorized into two domains, cognition-based EF, and emotion-related EF. The original concept of EF referred to cognition-based actions, and researchers and clinicians have used a variety of cognitive tests requiring deliberate attention to formally assess this type of EF. Such tests measure problem-solving, conceptual set-shifting, and rapid generation of verbal or nonverbal responses. Another form of action selection has been called emotion-related⁷⁹. Action selection at this level is based on rewards and punishments (positive and negative reinforcements) obtained in the past in similar situations. This emotion-related EF can be assessed using tests that measure the ability to modify behavior in response to changing reinforcement conditions⁸⁰. Anyway, FASD shows impairments for both domains, having them appear to be good predictors of behavioral problems in alcohol-affected people⁸¹⁻⁸⁴. Therefore, identifying and treating EF deficits overall years earlier may consistently help reduce the severity of problems including disruptive behavior and learning deficits^{85,86}.

In addition, a large body of neuropsychological research on children with FASD highlights a pattern where affected children succeed at simple tasks but display greater difficulty on more complex tasks⁸⁷. This illustrates how PAE disrupts the processing and integration of complex information, especially when they are required to hold and manipulate information in working memory⁸⁷.

GENERAL INTELLIGENCE

One of the first neurocognitive findings related to exposure to alcohol during pregnancy has been diminished intellectual capacity⁵². FASD is known as a

preventable cause of intellectual disability⁸⁸. Intelligence is a general mental ability for reasoning, planning, solving problems, thinking abstractly, comprehending complexity, and learning. Most studies that assessed the overall intelligence of individuals prenatally exposed to alcohol have used the Wechsler Intelligence Scales. It includes the WAIS, the WISC, and the WPPSI (Wechsler Preschool and Primary Scale of Intelligence) and is used for the assessment of intellectual ability across the lifespan⁸⁹. Fortunately, only a minority of individuals with FASD are intellectually disabled (defined as overall IQ score <70 and adaptive disability), while the majority have borderline cognitive functioning⁹⁰. As for the factorial indexes of the Wechsler Scales, the Working and Memory Index was the most adversely affected, with loss of attention, concentration, immediate memory, and mathematical skills. As for the Wechsler Scales subtests, there have been found more severe deficits in the arithmetic⁹¹. In a very recent study, a generally low performance of children with FASD was confirmed on all subtests of WISC, showing a significant weakness in working memory but also in processing speed⁹². IQ score has been found to have a weak correlation with adaptive functioning in this population^{93,94}. Furthermore, IQ score is significantly and inversely correlated with psychopathology. Therefore, children with moderate and severe intellectual disability experience greater psychiatric disturbance, and IQ scores below 50 indicate poor psychiatric outcomes⁹⁵.

Behavioral impairments

Behavioral problems of individuals with FASD are strictly connected with deficits in executive functioning resulting in difficulties in emotional regulation and adaptive functioning^{66,73,96}.

SELF-REGULATION

Self-regulation refers to the ability to control behavior and manage thoughts and emotions in appropriate ways. Unfortunately, individuals with FASD can exhibit a variety of dysregulation problems including disruptions in attention, problematic behavior, poor academic performance and adaptive functioning, and low social competence⁹⁷⁻⁹⁹. FASD has been found to exhibit serious internalizing (e.g. depression) and externalizing (e.g., hyperactivity, antisocial behavior) behavioral problems that often persist in adolescence and adulthood^{100,101}, high rates of mood disorders⁶¹ and oppositional defiant disorder, conduct disorder, and ADHD¹⁰². Importantly, ADHD is the most common psychiatric disorder diagnosed in children¹⁰³.

Behaviorally, children with FASDs have high also rates of social behavioral problems resulting from dif-

ficulties in social cognition and emotional processing¹⁰⁴. They have problems dealing with overstimulation and the frequent occurrence of externalizing behaviors¹⁰⁵. Furthermore, high rates of oppositional defiant disorder and conduct disorder often suggest severe difficulty following rules or controlling impulsive behaviors related to FASD^{5,106}. Interestingly, a variety of research is providing evidence that early interventions targeting self-regulation and executive functions among children with FASD produce more effective and generalizable improvements than domain-specific interventions¹⁰⁷⁻¹⁰⁹.

ADAPTIVE FUNCTIONING

PAE is often associated with a lower acquisition of life skills. Adaptive functioning, used to refer to skills related to daily living for personal and social sufficiency, is an area of special concern for children with FASD because these impairments are pervasive across domains and situations as children age¹¹⁰. There are three main areas of adaptive skills: social (e.g., communication, relationship development), practical (e.g., grooming, dressing, eating), conceptual (e.g., reading, writing), and motor skills for very young children¹¹¹. Deficits in adaptive functioning skills have been reported across the spectrum of FASD and in all three domains^{52,66,112}.

Discussion

FASD is considered a “hidden disability” because most individuals affected by PAE are not identified until adolescence or adulthood, if at all¹¹³⁻¹¹⁵. Misdiagnoses have major repercussions for treatment, which can be ineffective because the brain of FASDs works differently from similar neurodevelopmental diseases¹¹⁶. The current diagnostic FASD guidelines have the important merit of providing criteria for making a diagnosis, but they are concerned with defining the severity of the neurodevelopmental impairments rather than also the specificity of the impairments across the spectrum¹¹⁷.

This limit has impacted the diagnosis of ARND based primarily on the neurodevelopmental impairments rather than the characteristic facial traits and growth deficits associated only with FAS and pFAS. Unfortunately, 80-90% of cases of FASD are ARND representing a large FASD population that is often undiagnosed¹¹⁸. Since the cognitive and behavioral deficits seen in FASD are also common in other neurodevelopmental disorders, researchers have sought to identify a distinct neurobehavioral profile to facilitate the differential diagnosis of ARND. Nevertheless, all the studies on topic present significant methodological limitations, so that is not possible to define

a neurocognitive and behavioral profile specific to FASD¹¹⁹.

Conclusions

Preventing FASD begins with raising awareness among expectant mothers about the risks associated with alcohol consumption during pregnancy^{25,120,121}. Education campaigns emphasizing the importance of abstinence from alcohol throughout pregnancy play a crucial role in preventing FAS. It could be quite useful also to analyze, during pregnancy the presence of ethanol metabolites in the mother biological samples (hair, urine) or in the meconium to really disclose alcohol abuse during gestation^{24,122-127}. Healthcare providers also play a vital role by offering guidance and support to pregnant women, encouraging them to abstain from alcohol and providing resources for assistance if needed. Additionally, creating supportive environments that promote healthy lifestyle choices during pregnancy, coupled with policies that restrict access to alcohol among pregnant women, can contribute to the prevention of FAS and improve maternal and child health outcomes.

**Interdisciplinary Study Groups:* - Sapienza Università di Roma, AIDEFAD - Associazione Italiana Disordini da Esposizione Fetale ad Alcol e/o Droghe, SITAC - Società italiana per il trattamento dell'alcolismo e delle sue complicanze. SIFASD - Società Italiana Sindrome Feto-Alcolica, SIPPS - Società Italiana di Pediatria Preventiva e Sociale, FIMMG-Lazio - Federazione Italiana dei Medici di Medicina Generale Lazio, SIM-PeSV - Società Italiana di Medicina di Prevenzione e degli Stili di Vita, CIPE - Confederazione Italiana Pediatri. Alberto Chiriatti, Alberto Spalice, Alessio D'Angelo, Andrea Agostini, Andrea Liberti, Antonella Cavalieri, Camilla Di Dio, Cinzia Di Matteo, Elena Pacella, Enrico Finale, Francesco Chiarelli, Francesco D'Antonio, Giampiero Ferraguti, Ginevra Micangeli, Lina Corbi, Lucia Ruggieri, Maria Grazia Piccioni, Maria Pia Graziani, Mario Vitali, Martina Peracchini, Michela Menghi, Monica Napolitano, Paola Ciolli, Patrizia Riscica, Pier Luigi Bartoletti, Raffaella Punzo, Roberto Paparella, Romolo Di Iorio, Simona Vescina, Stefania Pipitone.

Conflict of interest: the authors have no conflict of interest to declare.

References

1. Terracina S, Ferraguti G, Tarani L, et al. Transgenerational abnormalities induced by paternal preconceptual alcohol drinking. Findings from humans and animal models. *Curr Neuropsychopharmacol* 2021; 19: 1158-73.
2. Basavarajappa BS, Subbanna S. Synaptic plasticity abnormalities in Fetal Alcohol Spectrum Disorders. *Cells* 2023; 12: 442.
3. Vorgias D, Bernstein B. Fetal Alcohol Syndrome. In: *Treasure Island (FL), Treasure Island (FL)*, 2021.
4. Hoyme HE, Kalberg WO, Elliott AJ, et al. Updated clinical guidelines for diagnosing fetal alcohol spectrum disorders. *Pediatrics* 2016; 138: e20154256-e20154256.
5. Mattson SN, Bernes GA, Doyle LR. Fetal Alcohol Spectrum Disorders: a review of the neurobehavioral deficits associated with prenatal alcohol exposure. *Alcohol Clin Exp Res* 2019; 43: 1046-62.
6. Morton J, Frith U. Causal modelling a structural approach to developmental psychopathology. In: Cicchetti D, Cohen DJ (eds). *Developmental psychopathology*. Vol. 1. Theory and methods. Hoboken: NJ: John Wiley & Sons, 1995.
7. Rockhold M, Donald K, Kautz-Turnbull C, Petrenko C. Neuroimaging findings in FASD across the lifespan. In: Abdul-Rahman OA, Petrenko CLM (eds). *Fetal Alcohol Spectrum Disorders*. Cham, CH: Springer International Publishing, 2023.
8. Boateng T, Beauchamp K, Torres F, et al. Brain structural differences in children with fetal alcohol spectrum disorder and its subtypes. *Front Neurosci* 2023; 17: 1152038.
9. Roebuck-Spencer TM, Mattson SN, Marion SD, Brown WS, Riley EP. Bimanual coordination in alcohol-exposed children: role of the corpus callosum. *J Int Neuropsychol Soc* 2004; 10: 536-48.
10. Coles CD, Platzman KA, Lynch ME, Freides D. Auditory and visual sustained attention in adolescents prenatally exposed to alcohol. *Alcohol Clin Exp Res* 2002; 26: 263-71.
11. Sowell ER, Mattson SN, Thompson PM, Jernigan TL, Riley EP, Toga AW. Mapping callosal morphology and cognitive correlates: Effects of heavy prenatal alcohol exposure. *Neurology* 2001; 57: 235-44.
12. Gautam P, Nuñez SC, Narr KL, Kan EC, Sowell ER. Effects of prenatal alcohol exposure on the development of white matter volume and change in executive function. *NeuroImage Clin* 2014; 5: 19-27.
13. Mattson SN, Riley EP, Sowell ER, Jernigan TL, Sobel DF, Jones KL. A decrease in the size of the basal ganglia in children with fetal alcohol syndrome. *Alcohol Clin Exp Res* 1996; 20: 1088-93.
14. Astley SJ, Aylward EH, Olson HC, et al. Magnetic resonance imaging outcomes from a comprehensive magnetic resonance study of children with fetal alcohol spectrum disorders. *Alcohol Clin Exp Res* 2009; 33: 1671-89.
15. Inkelis SM, Moore EM, Bischoff-Grethe A, Riley EP. Neurodevelopment in adolescents and adults with fetal alcohol spectrum disorders (FASD): a magnetic resonance region of interest analysis. *Brain Res* 2020; 1732.
16. Treit S, Chen Z, Zhou D, et al. Sexual dimorphism of volume reduction but not cognitive deficit in fetal alcohol spectrum disorders: a combined diffusion tensor imaging, cortical thickness and brain volume study. *NeuroImage Clin* 2017; 15: 284-97.
17. Chen ML, Olson HC, Picciano JF, Starr JR, Owens J. Sleep problems in children with fetal alcohol spectrum disorders. *J Clin Sleep Med* 2012; 8: 421-9.
18. Meombe Mbolle A, Thapa S, Bukiya AN, Jiang H. High-resolution imaging in studies of alcohol effect on prenatal development. *Adv Drug Alcohol Res* 2023; 3: 10790.
19. Wozniak JR, Mueller BA, Mattson SN, et al. Functional connectivity abnormalities and associated cognitive deficits in fetal alcohol Spectrum disorders (FASD). *Brain Imaging Behav* 2017; 11: 1432-45.
20. Lees B, Mewton L, Jacobus J, et al. Association of prenatal alcohol exposure with psychological, behavioral, and neurodevelopmental outcomes in children from the adolescent brain cognitive development study. *Am J Psychiatry* 2020; 177: 1060-72.
21. Sundermann B, Feldmann R, Mathys C, et al. Functional connectivity of cognition-related brain networks in adults with fetal alcohol syndrome. *BMC Med* 2023; 21: 496.
22. Terracina S, Tarani L, Ceccanti M, et al. The impact of oxidative stress on the epigenetics of Fetal Alcohol Spectrum Disorders. *Antioxidants* 2024; 13: 410.

23. Coriale G, Fiorentino D, Lauro FDI, et al. Fetal Alcohol Spectrum Disorder (FASD): neurobehavioral profile, indications for diagnosis and treatment. *Riv Psichiatr* 2013; 48: 359-69.
24. Coriale G, Ceccanti M, Fiore M, et al. Delay in the fine-tuning of locomotion in infants with meconium positive to biomarkers of alcohol exposure: a pilot study. *Riv Psichiatr* 2024; 59: 52-9.
25. Messina MP, D'Angelo A, Battagliese G, et al. Fetal alcohol spectrum disorders awareness in health professionals: implications for psychiatry. *Riv Psichiatr* 2020; 55: 79-89.
26. Aragón AS, Coriale G, Fiorentino D, et al. Neuropsychological characteristics of Italian children with fetal alcohol spectrum disorders. *Alcohol Clin Exp Res* 2008; 32: 1909-19.
27. Coriale G, Fiorentino D, Kodituwakku PW, et al. Identification of children with prenatal alcohol exposure. *Curr Dev Disord Reports* 2014; 1: 141-8.
28. Lezak MD, Howieson DB, Bigler ED, Tranel D. Neuropsychological assessment, 5th ed. Neuropsychological assessment. USA: Oxford University Press, 2004.
29. Gyllencreutz E, Aring E, Landgren V, Landgren M, Grönlund MA. Visual perception problems and quality of life in young adults with foetal alcohol spectrum disorders. *Acta Ophthalmol* 2022; 100: e115-21.
30. Lyubasyuk V, Jones KL, Caesar MA, Chambers C. Vision outcomes in children with fetal alcohol spectrum disorders. *Birth defects Res* 2023; 115: 1208-15.
31. Ayoub L, Aring E, Gyllencreutz E, et al. Visual and ocular findings in children with fetal alcohol spectrum disorders (FASD): validating the FASD Eye Code in a clinical setting. *BMJ Open Ophthalmol* 2023; 8: e001215.
32. Pennell J. Assessing and managing sensory processing. In: Mukherjee RAS, Aiton N (eds). *Prevention, recognition and management of Fetal Alcohol Spectrum Disorders*. Cham, CH: Springer International Publishing, 2021.
33. Fjeldsted B, Xue L. Sensory processing in young children with Fetal Alcohol Spectrum Disorder. *Phys Occup Ther Pediatr* 2019; 39: 553-65.
34. Bjuland KJ, Løhaugen GCC, Martinussen M, Skranes J. Cortical thickness and cognition in very-low-birth-weight late teenagers. *Early Hum Dev* 2013; 89: 371-80.
35. Johnson S. Cognitive and behavioural outcomes following very preterm birth. *Semin Fetal Neonatal Med* 2007; 12: 363-73.
36. Koenen KC, Moffitt TE, Poulton R, Martin J, Caspi A. Early childhood factors associated with the development of post-traumatic stress disorder: results from a longitudinal birth cohort. *Psychol Med* 2007; 37: 181-92.
37. Johnston D, Branton E, Rasmuson L, Schell S, Gross DP, Pritchard-Wiart L. Accuracy of motor assessment in the diagnosis of fetal alcohol spectrum disorder. *BMC Pediatr* 2019; 19: 171.
38. Moss N, Moss-Racusin L. Visual-motor coordination. In: *Practical guide to child and adolescent psychological testing. Best practices in child and adolescent behavioral health care*. Cham, CH: Springer International Publishing, 2021.
39. Lu H. A meta-analysis of handwriting and visual-motor integration. 2023. *Learn Individ Differ* 2024; 109: 102404.
40. Cook JL, Green CR, Lilley CM, et al.; Canada Fetal Alcohol Spectrum Disorder Research Network. Fetal alcohol spectrum disorder: a guideline for diagnosis across the lifespan. *CMAJ* 2016; 188: 191-7.
41. Doney R, Lucas BR, Jones T, Howat P, Sauer K, Elliott EJ. Fine motor skills in children with prenatal alcohol exposure or fetal alcohol spectrum disorder. *J Dev Behav Pediatr* 2014; 35: 598-609.
42. Doney R, Lucas BR, Watkins RE, et al. Visual-motor integration, visual perception, and fine motor coordination in a population of children with high levels of Fetal Alcohol Spectrum Disorder. *Res Dev Disabil* 2016; 55: 346-57.
43. Carsone B, Green K, Torrence W, Henry B. Systematic review of visual motor integration in children with developmental disabilities. *Occup Ther Int* 2021; 2021: 1801196.
44. Johnston D, Pritchard-Wiart L, Branton E, Gross D, Thompson-Hodgetts S. Pattern of visual-motor integration, visual perception, and fine motor coordination abilities in children being assessed for Fetal Alcohol Spectrum Disorder. *J Dev Behav Pediatr* 2023; 44: e463-e469.
45. Nath S, Szűcs D. Construction play and cognitive skills associated with the development of mathematical abilities in 7-year-old children. *Learn Instr* 2014; 32: 73-80.
46. Pei J, Denys K, Hughes J, Rasmussen C. Mental health issues in fetal alcohol spectrum disorder. *J Ment Heal* 2011; 20: 473-83.
47. Nesayan A, Amani M, Asadi Gandomani R. Cognitive profile of children and its relationship with academic performance. *Basic Clin Neurosci* 2019; 10: 165-74.
48. Castillo Castejón O, González I, Prieto E, Pérez T, Pablo LE, Pueyo V. Visual cognitive impairments in children at risk of prenatal alcohol exposure. *Acta Paediatr* 2019; 108: 2222-8.
49. Khatib L, Li Y, Geary D, Popov V. Meta-analysis on the relation between visuomotor integration and academic achievement: role of educational stage and disability. *Educ Res Rev* 2022; 35: 100412.
50. Wang L, Wang L. Relationships between motor skills and academic achievement in school-aged children and adolescents: a systematic review. *Children* 2024; 11: 336.
51. Sridhar S, Khamaj A, Asthana MK. Cognitive neuroscience perspective on memory: overview and summary. *Front Hum Neurosci* 2023; 17: 1217093.
52. Crocker N, Vaurio L, Riley EP, Mattson SN. Comparison of verbal learning and memory in children with heavy prenatal alcohol exposure or attention-deficit/hyperactivity disorder. *Alcohol Clin Exp Res* 2011; 35: 1114-21.
53. Mattson SN, Riley EP, Delis DC, Stern C, Jones KL. Verbal learning and memory in children with fetal alcohol syndrome. *Alcohol Clin Exp Res* 1996; 20: 810-6.
54. Roediger DJ, Krueger AM, de Water E, et al. Hippocampal subfield abnormalities and memory functioning in children with fetal alcohol spectrum disorders. *Neurotoxicol Teratol* 2021; 83: 106944.
55. Mattson S, Roebuck-Spencer T. Acquisition and retention of verbal and nonverbal information in children with heavy prenatal alcohol exposure. *Alcohol Clin Exp Res* 2002; 26: 875-82.
56. Pei JR, Rinaldi CM, Rasmussen C, Massey V, Massey D. Memory patterns of acquisition and retention of verbal and nonverbal information in children with fetal alcohol spectrum disorders. *Can J Clin Pharmacol* 2008; 15: e44-56.
57. Willford J, Richardson G, Leech S, Day N. Verbal and visuospatial learning and memory function in children with moderate prenatal alcohol exposure. *Alcohol Clin Exp Res* 2004; 28: 497-507.
58. Kodituwakku P, Coriale G, Fiorentino D, et al. Neurobehavioral characteristics of children with fetal alcohol spectrum disorders in communities from Italy: preliminary results. *Alcohol Clin Exp Res* 2006; 30: 1551-61.
59. Green CR, Mihic AM, Nikkel SM, et al. Executive function deficits in children with fetal alcohol spectrum disorders (FASD) measured using the Cambridge Neuropsychological Tests Automated Battery (CANTAB). *J Child Psychol Psychiatry Allied Discip* 2009; 50: 688-97.
60. Hendricks G, Malcolm-Smith S, Adnams C, Stein DJ, Donald KAM. Effects of prenatal alcohol exposure on language, speech and communication outcomes: a review longitudinal studies. *Acta Neuropsychiatr* 2019; 31: 74-83.

61. O'Connor MJ, Paley B. The relationship of prenatal alcohol exposure and the postnatal environment to child depressive symptoms. *J Pediatr Psychol* 2006; 3: 50-64.
62. Gray R, Gaska J, Winterbottom M. Relationship between sustained, orientated, divided, and selective attention and simulated aviation performance: training & pressure effects. *J Appl Res Mem Cogn* 2016; 5: 34-42.
63. Moisa M, Salmela V, Hietajärvi L, et al. Media multitasking is associated with distractibility and increased prefrontal activity in adolescents and young adults. *Neuroimage* 2016; 134: 113-21.
64. Mattson JT, Thorne JC, Kover ST. Relationship between task-based and parent report-based measures of attention and executive function in children with Fetal Alcohol Spectrum Disorders (FASD). *J Pediatr Neuropsychol* 2020; 6: 176-88.
65. Coles C. Fetal alcohol exposure and attention: moving beyond ADHD. *Alcohol Res Health* 2001; 25: 199-203.
66. Popova S, Charness ME, Burd L, et al. Fetal alcohol spectrum disorders. *Nat Rev Dis Prim* 2023; 9: 11.
67. Mattson SN, Calarco KE, Lang AR. Focused and shifting attention in children with heavy prenatal alcohol exposure. *Neuropsychology* 2006; 20: 361-9.
68. Coles CD, Platzman KA, Raskind-Hood CL, Brown RT, Falek A, Smith IE. A comparison of children affected by prenatal alcohol exposure and attention deficit, hyperactivity disorder. *Alcohol Clin Exp Res* 1997; 21: 150-61.
69. Cristofori I, Cohen-Zimmerman S, Grafman J. Executive functions. *Handb Clin Neurol* 2019; 163: 197-219.
70. D'Acremont M, der Linden M. Adolescent impulsivity: findings from a community sample. *J Youth Adolesc* 2005; 34: 427-35.
71. Cristofori I, Cohen-Zimmerman S, Grafman J. Executive functions. *Handb Clin Neurol* 2019; 163: 197-219.
72. Fuglestad AJ, Whitley ML, Carlson SM, et al. Executive functioning deficits in preschool children with Fetal Alcohol Spectrum Disorders. *Child Neuropsychol* 2015; 21: 716-31.
73. Kodituwakku PW. Defining the behavioral phenotype in children with fetal alcohol spectrum disorders: a review. *Neurosci Biobehav Rev* 2007; 31: 192-201.
74. Nash K, Stevens S, Greenbaum R, Weiner J, Koren G, Rovet J. Improving executive functioning in children with fetal alcohol spectrum disorders. *Child Neuropsychol* 2015; 21: 191-209.
75. Coles CD, Platzman KA, Raskind-Hood CL, Brown RT, Falek A, Smith IE. A comparison of children affected by prenatal alcohol exposure and attention deficit, hyperactivity disorder. *Alcohol Clin Exp Res* 1997; 21: 150-61.
76. Kingdon D, Cardoso C, McGrath JJ. Research review: executive function deficits in fetal alcohol spectrum disorders and attention-deficit/hyperactivity disorder - a meta-analysis. *J Child Psychol Psychiatry Allied Discip* 2016; 57: 116-31.
77. Mattson SN, Goodman AM, Caine C, Delis DC, Riley EP. Executive functioning in children with heavy prenatal alcohol exposure. *Alcohol Clin Exp Res* 1999; 23: 1808-15.
78. Fuglestad AJ, Whitley ML, Carlson SM, et al. Executive functioning deficits in preschool children with Fetal Alcohol Spectrum Disorders. *Child Neuropsychol* 2015; 21: 716-31.
79. Rolls ET, Treves A. Neural networks in the brain involved in memory and recall. In: Van Pelt J, Corner MA, Uylings HBM, Lopes Da Silva FH (eds). *the self-organizing brain: from growth cones to functional networks*. Amsterdam: Elsevier, 1994.
80. Kodituwakku PW, Kalberg W, May PA. The effects of prenatal alcohol exposure on executive functioning. *Alcohol Res Health* 2001; 25: 192-8.
81. Coriale G, Gencarelli S, Battagliese G, et al. Physiological responses to induced stress in individuals affected by alcohol use disorder with dual diagnosis and alexithymia. *Clin Ter* 2020; 171: e120-9.
82. D'Angelo A, Peracchini M, Agostini A, et al. the impact of oxidative stress on pregnancy. The Neglected role of alcohol misuse. *Clin Ter* 2024; 175: 47-56.
83. D'Angelo A, Petrella C, Greco A, et al. Acute alcohol intoxication: a clinical overview. *Clin Ter* 2022; 173: 280-91.
84. Profeta G, Micangeli G, Tarani F, et al. Sexual developmental disorders in pediatrics. *Clin Ter* 2022; 173: 475-88.
85. Betts JL, Eggins E, Chandler-Mather N, et al. Interventions for improving executive functions in children with foetal alcohol spectrum disorder (FASD): a systematic review. *Campbell Syst Rev* 2022; 18: e1258.
86. Christine Christensen Løhaugen G, Cecilie Tveiten A, Skranes J. Interventions for children and adolescents with Fetal Alcohol Spectrum Disorders (FASD). In: Patel VB, Preedy VR (eds). *Handbook of substance misuse and addictions*. Cham, CH: Springer International Publishing, 2022.
87. Kodituwakku PW. Defining the behavioral phenotype in children with fetal alcohol spectrum disorders: a review. *Neurosci Biobehav Rev* 2007; 31: 192-201.
88. Williams JF, Smith VC. Fetal Alcohol Spectrum Disorders. *Pediatrics* 2015; 136: e1395-406.
89. Goldstein G, Mazefsky C. Wechsler scales of intelligence. In: Volkmar FR (ed). *Encyclopedia of Autism Spectrum Disorders*. New York, NY: Springer New York, 2013.
90. Mattson SN, Riley EP. Parent ratings of behavior in children with heavy prenatal alcohol exposure and IQ-matched controls. *Alcohol Clin Exp Res* 2000; 24: 226-31.
91. Nash K, Sheard E, Rovet J, Koren G. Understanding fetal alcohol spectrum disorders (FASDs): toward identification of a behavioral phenotype. *ScientificWorldJournal* 2008; 8: 873-82.
92. Kerdreux E, Fraize J, Garzón P, et al. Questioning cognitive heterogeneity and intellectual functioning in fetal alcohol spectrum disorders from the Wechsler intelligence scale for children. *Clin Neuropsychol* 2024; 38: 1109-1132.
93. Boseck J, Davis A, Cassady J, Finch H, Gelder B. Cognitive and adaptive skill profile differences in children with attention-deficit hyperactivity disorder with and without comorbid Fetal Alcohol Spectrum Disorder. *Appl Neuropsychol Child* 2014; 4: 1-7.
94. Kautz-Turnbull C, Petrenko CLM. A meta-analytic review of adaptive functioning in fetal alcohol spectrum disorders, and the effect of IQ, executive functioning, and age. *Alcohol Clin Exp Res* 2021; 45: 2430-47.
95. Spohr HL, Willms J, Steinhausen HC. The fetal alcohol syndrome in adolescence. *Acta Paediatr Suppl* 1994; 404: 19-26.
96. Lange S, Shield K, Rehm J, Anagnostou E, Popova S. Fetal alcohol spectrum disorder: neurodevelopmentally and behaviorally indistinguishable from other neurodevelopmental disorders. *BMC Psychiatry* 2019; 19: 322.
97. Brown JM, Trnka A, Harr D, Dodson KD, Wartnik HAP, Donaldson K. Fetal alcohol spectrum disorder (FASD): a beginner's guide for mental health professionals. *J Neurol Clin Neurosci* 2018; 2: 13-9.
98. Mattson SN, Crocker N, Nguyen TT. Fetal alcohol spectrum disorders: neuropsychological and behavioral features. *Neuropsychol Rev* 2011; 21: 81-101.
99. Gill K, Thompson-Hodgetts S. Self-regulation in fetal alcohol spectrum disorder: a concept analysis. *J Occup Ther Sch Early Interv* 2018; 11: 329-45.
100. Barr HM, Bookstein FL, O'Malley KD, Connor PD, Huggins JE, Streissguth AP. Binge drinking during pregnancy as a predictor of psychiatric disorders on the Structured

- Clinical Interview for DSM-IV in young adult offspring. *Am J Psychiatr* 2006; 163: 1061-5.
101. Tsang TW, Lucas BR, Carmichael Olson H, Pinto RZ, Elliott EJ. Prenatal alcohol exposure, FASD, and child behavior: a meta-analysis. *Pediatrics* 2016; 137: e20152542.
 102. Kapasi A, Pei J, Joly V, et al. Exploring Self-Regulation Strategy Use in Adolescents with FASD. *Journal of Occupational Therapy, Schools, & Early Intervention* 2021; 14: 184-206.
 103. Young S, Absoud M, Blackburn C, et al. Guidelines for identification and treatment of individuals with attention deficit/hyperactivity disorder and associated fetal alcohol spectrum disorders based upon expert consensus. *BMC Psychiatry* 2016; 16: 324.
 104. Greenbaum RL, Stevens SA, Nash K, Koren G, Rovet J. Social cognitive and emotion processing abilities of children with fetal alcohol spectrum disorders: a comparison with attention deficit hyperactivity disorder. *Alcohol Clin Exp Res* 2009; 33: 1656-70.
 105. Peadar E. Distinguishing between attention-deficit hyperactivity and fetal alcohol spectrum disorders in children: clinical guidelines. *Neuropsychiatr Dis Treat* 2010; 6: 509.
 106. Doyle LR, Glass L, Wozniak JR, et al. Relation between oppositional/conduct behaviors and executive function among youth with histories of heavy prenatal alcohol exposure. *Alcohol Clin Exp Res* 2019; 43: 1135-44.
 107. Kodituwakku PW. A neurodevelopmental framework for the development of interventions for children with fetal alcohol spectrum disorders. *Alcohol* 2010; 44: 717-28.
 108. Kable JA, Taddeo E, Strickland D, Coles CD. Improving FASD children's self-regulation: piloting Phase 1 of the GoFAR intervention. *Child Fam Behav Ther* 2016; 38: 124-41.
 109. Betts JL, Eggins E, Chandler-Mather N, et al. Interventions for improving executive functions in children with foetal alcohol spectrum disorder (FASD): a systematic review. *Campbell Syst Rev* 2022; 18: e1258.
 110. Streissguth AP, Aase JM, Clarren SK, Randels SP, LaDue RA, Smith DF. Fetal Alcohol Syndrome in Adolescents and Adults. *JAMA* 1991; 265: 1961.
 111. Sparrow SS, Cicchetti DV. Diagnostic uses of the vine-land adaptive behavior scales. *J Pediatr Psychol* 1985; 10: 215-25.
 112. Crawford A, Te Nahu LTH, Peterson ER, McGinn V, Robertshaw K, Tippet L. Cognitive and social/emotional influences on adaptive functioning in children with FASD: clinical and cultural considerations. *Child Neuropsychol* 2020; 26: 1112-44.
 113. Ceccanti M, Hamilton D, Coriale G, et al. Spatial learning in men undergoing alcohol detoxification. *Physiol Behav* 2015; 149: 324-30.
 114. Ceccanti M, Coriale G, Hamilton DA, et al. Virtual Morris task responses in individuals in an abstinence phase from alcohol. *Can J Physiol Pharmacol* 2018; 96: 128-36.
 115. Streissguth AP, Bookstein FL, Barr HM, Sampson PD, O'Malley K, Young JK. Risk factors for adverse life outcomes in Fetal Alcohol Syndrome and Fetal Alcohol Effects. *J Dev Behav Pediatr* 2004; 25: 228-38.
 116. Chasnoff IJ, Wells AM, King L. Misdiagnosis and missed diagnoses in foster and adopted children with prenatal alcohol exposure. *Pediatrics* 2015; 135: 264-70.
 117. Lange S, Probst C, Gmel G, Rehm J, Burd L, Popova S. Global prevalence of fetal alcohol spectrum disorder among children and youth: A systematic review and meta-analysis. *JAMA Pediatr* 2017; 171: 948-56.
 118. Chudley AE. Fetal alcohol spectrum disorder: counting the invisible - mission impossible? *Arch Dis Child* 2008; 93: 721-2.
 119. Maya-Enero S, Ramis-Fernández SM, Astals-Vizcaino M, García-Algar Ó. Neurocognitive and behavioral profile of fetal alcohol spectrum disorder. *An Pediatr* 2021; 95: 208.e1-208.e9.
 120. Messina MP, D'Angelo A, Ciccarelli R, et al. Knowledge and practice towards alcohol consumption in a sample of university students. *Int J Environ Res Public Health* 2021; 18: 9528.
 121. Messina MP, D'Angelo A, Battagliese G, et al. Fetal alcohol spectrum disorders awareness in health professionals: implications for psychiatry. *Riv Psichiatr* 2020; 55: 79-89.
 122. Mattia A, Moschella C, David MC, et al. Development and Validation of a GC-EI-MS/MS Method for Ethyl Glucuronide Quantification in Human Hair. *Front Chem* 2022; 10: 858205.
 123. Ceci FM, Fiore M, Agostinelli E, et al. Urinary ethyl glucuronide for the assessment of alcohol consumption during pregnancy: comparison between biochemical data and screening questionnaires. *Curr Med Chem* 2021; 29: 3125-41.
 124. Ferraguti G, Ciolli P, Carito V, et al. Ethylglucuronide in the urine as a marker of alcohol consumption during pregnancy: comparison with four alcohol screening questionnaires. *Toxicol Lett* 2017; 275: 49-56.
 125. Gomez-Roig MD, Marchei E, Sabra S, et al. Maternal hair testing to disclose self-misreporting in drinking and smoking behavior during pregnancy. *Alcohol* 2018; 67: 1-6.
 126. Morini L, Marchei E, Tarani L, et al. Testing ethylglucuronide in maternal hair and nails for the assessment of fetal exposure to alcohol: Comparison with meconium testing. *Ther Drug Monit* 2013; 35: 402-7.
 127. Ferraguti G, Merlino L, Battagliese G, et al. Fetus morphology changes by second-trimester ultrasound in pregnant women drinking alcohol. *Addict Biol* 2020; 25: e12724.