

Psychotherapeutic treatment of eating disorders improve dissociative experiences and impulse regulation but not alexithymia. A case series report

Il trattamento psicoterapeutico dei disturbi alimentari migliora le tendenze dissociative e impulsive, ma non l'alestitimia.

Una serie di casi clinici

MANUELA CASLINI^{1,2}, LAURA RIVOLTA^{2,3}, LUIGI ENRICO ZAPPA², GIUSEPPE CARRÀ⁴,
MASSIMO CLERICI^{2,5}

E-mail: m.caslini3@campus.unimib.it

¹PhD School in Neuroscience, Department of Surgery and Translational Medicine, University of Milano Bicocca, Monza, MB, Italy

²Specialist service for eating disorders, Psychiatry Operating Unit, San Gerardo Hospital Care Trust, Monza, MB, Italy

³Department of Psychology, University of Milano Bicocca, Milan, Italy.

⁴Division of Psychiatry, Faculty of Brain Sciences, University College London, United Kingdom

⁵Department of Surgery and Translational Medicine, University of Milano Bicocca, Monza, MB, Italy

SUMMARY. Aim. Eating disorders (EDs) are complex conditions associated with disability and a high rate of mortality. Typical characteristics of these diseases are dissociation, alexithymia and impulse dysregulation, all strategies dealing with negative emotions and regulate negative affect and anxiety. Our study aimed to assess the effectiveness of intensive psychological treatment for EDs, with particular reference to the above mentioned clinical characteristics. **Methods.** Eight outpatients with eating disorders in psychotherapeutic treatment were evaluated in two stages after one year (T1 and T2), using the Eating Disorder Inventory II, the Toronto Alexithymia Scale 20, and the Dissociative Experiences Scale. **Results.** Wilcoxon test showed significant reductions in DES score as well as in two subscales of the EDI-2, Impulse Regulation and Body Dissatisfaction (I-EDI2 and BD-EDI2), while alexithymia levels did not show any difference. **Conclusions.** We can confirm the effectiveness of psychotherapy in people with EDs as regards dissociative moments, impulsivity and body dissatisfaction. However, alexithymia remains unchanged, possibly because of its deep emotional nature.

KEY WORDS: eating disorder, psychotherapy, dissociation, impulse regulation, alexithymia.

RIASSUNTO. Scopo. I disturbi alimentari (DCA) sono condizioni complesse associate a importanti disabilità e a un alto tasso di mortalità. In queste condizioni psicopatologiche si possono riscontrare alestitimia, tendenze dissociative e impulsive, messe in atto quali strategie di gestione delle emozioni negative e di regolazione affettiva. Questo studio si propone di valutare l'efficacia di un trattamento psicologico intensivo per DCA, con particolare riferimento alle caratteristiche cliniche citate. **Metodi.** Otto pazienti con DCA seguite in trattamento psicoterapico in un programma ambulatoriale multidisciplinare sono state valutate in due tempi (T1 e T2), a distanza di un anno, mediante i questionari autosomministrati Eating Disorder Inventory II (EDI-2), Toronto Alexithymia Scale-20 (TAS-20) e Dissociative Experiences Scale (DES). **Risultati.** Il test di Wilcoxon ha mostrato una significativa riduzione nel punteggio DES e in due sotto-scale EDI-2 - Regolazione degli impulsi e Insoddisfazione Corporea (I-EDI2 e BD-EDI2). Nei punteggi della TAS-20, al contrario, non è stato riscontrato alcun cambiamento. **Conclusioni.** I risultati confermano l'efficacia della psicoterapia rispetto ai momenti dissociativi, alla regolazione degli impulsi e all'insoddisfazione corporea. Tuttavia, forse a causa della sua profonda natura emotiva, l'alestitimia rimane stabile.

PAROLE CHIAVE: disturbi del comportamento alimentare, psicoterapia, dissociazione, regolazione degli impulsi, alestitimia.

INTRODUCTION

Eating disorders (EDs) are complex conditions, with a common theme on food/weight obsession, associated with disability and the highest rates of mortality rates among psychiatric disorders¹. Lifetime prevalence estimates among women of anorexia nervosa (AN), bulimia nervosa (BN) and

binge eating disorder (BED) are 0.9%, 1.5%, and 3.5%², respectively.

Given the increasing prevalence of these disorders, to date it is unclear why the disease develops in some people and not in others. Definition of the specific psychological variables intervening in the disease process would be essential.

Dissociation, alexithymia and impulse regulation are associated with EDs in order to deal with negative emotions and regulate negative affect and anxiety³, making all of them appropriate treatment targets.

Dissociation – «a disruption in the integrated functions of consciousness, memory, identity, and perception of the environment»⁴ – is frequent among patients with EDs^{5,6}, and often associated with childhood abuse or traumas⁷⁻⁹. Dissociation seems important in determining bingeing severity¹⁰⁻¹², and relevant amnesia and timelessness, as well as feelings of depersonalization and derealisation¹³, are considered common properties of a binge episode¹⁴. Emotional disavowal, disturbed body image and deficits in the ongoing development of Self Concept may link EDs with dissociation^{15,16} but also with self-injurious behaviours¹⁷, similarly typical features of the disorder.

In addition alexithymia, with its four clusters of cognitive and affective characteristics¹⁸ and assessed by the Toronto Alexithymia Scale (TAS20)¹⁹, has been reported in people with EDs with rates between 23 and 77%²⁰. Patients with EDs indeed show significant difficulties in identifying their feelings and expressing them verbally, attempting to avoid feared sensations and emotional experiences in general, with reduced ability to symbolize²¹, partly explaining poor response to treatment²².

Finally, impulsivity, a behavioural aspect of emotion dysregulation²³, has been found to be associated with EDs characterized by bingeing and/or purging²⁴ regulating negative affect and anxiety.

Most people with eating disorders should be managed on an outpatient basis with psychological treatment mainly focussed on peculiar cognitive, emotional, and behavioural and personality attitudes such as alexithymia an affect dysregulation, impulse discontrol and dissociation proneness²⁵. In the absence of robust evidence, cognitive behavior therapy has shown preliminary, promising results²⁶, but there is the need for further experimental research before recommending that for implementation. Our study aimed to assess the effectiveness of intensive psychological treatment for EDs, with particular reference to dissociation, alexithymia and impulse regulation.

METHODS

Setting

The research project was approved by the Ethics Committee of Milano Bicocca University, and carried out at the Specialist Practice for EDs, San Gerardo Hospital Care Trust, Monza, Italy.

Treatment model

After a psychiatric assessment followed by a multidisciplinary evaluation, including physical condition, three female psychotherapists, with ten years of experience in treating ED patients, delivered the treatment. This was based on cognitive restructuring techniques, emotional literacy and regulation, aiming at reducing ED symptoms: weight recovery and/or cessation of binge eating and food dysregulation, but also dissociation proneness, alexithymia, impulse regulation, pathological perfection-

ism, brooding, distorted body image, control and sense of personal worthlessness²⁷. This was complemented with dynamic-oriented, supportive psychotherapy elements, building a bond of trust that makes possible to broaden the self-perception and the self-organization with the reduction of the pathological split between body and mind, and identifying emotions and thoughts, regenerating connections with the experience of an affective state collapsed in eating symptom²⁸. The psychotherapeutic treatment, generally, can last more than a year with one session per week.

Participants and procedures

The sample consisted of eight consecutively admitted female subjects with EDs. Axis I and II diagnostic evaluation was made according to DSM IV-TR criteria²⁹ and personality was assessed with the Q-sort SWAP-200³⁰.

Baseline and one year follow-up assessments were completed on January and December 2011 with the following set of tools.

Eating Disorders Inventory 2

This 91-item self-report tool measures various symptoms of eating disorders^{31,32}. It also provides standardized subscale scores on 11 clinically relevant dimensions, i.e., Drive for thinness (DT), Bulimia (BU), Body dissatisfaction (BD), Ineffectiveness (IN), Perfectionism (P), Interpersonal distrust (ID), Interoceptive awareness (IA), Maturity fears (MF), Asceticism (A), Impulse regulation (I), Social insecurity (SI). In the present study, for people with anorexia nervosa, we considered as a measure of efficacy also the BMI.

The Dissociative Experiences Scale

The Dissociative Experiences Scale (DES) is a widely used self-report instrument that measures dissociation³³. It includes 28 questions on various dissociative experiences. For each item, respondents score between 0 and 100% and the general score is calculated by the average of all the answers. A DES cut-off score of 15 to 20 is characteristic of dissociative disorders³⁴. In the present study we treated DES as a continuous variable.

Toronto Alexithymia Scale

TAS-20 is the most widely used self-report measure of alexithymia³⁵⁻³⁷. It consists of three subscales, difficulty in identifying feelings and distinguishing them from bodily sensation (DIF); difficulty in describing feelings to others (DDF); and an externally oriented style of thinking (EOT). Respondents use for each item a Likert scale ranging from 1 (strongly disagree) to 5 (strongly agree).

Data analysis

We used Statistics 6 for Windows for data processing. First, normal distribution of the variables investigated – DES, TAS20, EDI 2 – was tested with Kolmogorov-Smirnov test. In order to evaluate the improvement in dissociative experiences, alexithymia and impulse regulation, we carried out relevant pre-post comparisons (mean and SD) using Wilcoxon test. Statistical significance level was set at $p < .05$.

Psychotherapeutic treatment of eating disorders may improve dissociative experiences and impulse regulation but not alexithymia

RESULTS

Descriptive data of individuals in the clinical sample are shown in Table 1.

The sample comprised eight middle class female subjects, aged between 19 and 45 years, Italian native, with 13.12 (SD=2.47) years of education. According to DSM-IV-TR criteria, three subjects were suffering from restrictive anorexia nervosa (R-AN), two from bulimia nervosa (BN), and three from eating disorders not otherwise specified (EDNOS). Not surprisingly³⁸, more than half of the patients met criteria for an Axis II diagnosis and had had previous treatments.

Six out of eight subjects had a score >50 on the SWAP200 – High Functioning Scale³⁰, with a good overall functioning making them appropriate for psychological treatment. Furthermore, we built a checklist, including all core characteristics typical of EDs, regardless of single disorder’s features (Table 2). All 8 patients met listed characteristics.

Kolmogorov-Smirnov test showed non-normal distribution for all the three variables – DES, TAS20, EDI 2 – thus, non-parametrical statistical tests have been used.

One year follow-up

We tested pre- post-treatment (T1/T2) differences in the experimental sample. Wilcoxon test showed significant reductions in DES score as well as in two subscales of the EDI-2, Impulse Regulation and Body Dissatisfaction (I-EDI2 e BD-EDI2). However, as measured by TAS-20, alexithymia did not show any change (Table 3). Furthermore the three subjects with Anorexia Nervosa showed a stable (one) and higher (two) BMI (Table 1).

DISCUSSION

We could study a wide range of women with EDs, sharing core EDs characteristics, regardless of single disorder’s cate-

gorical features. Our study shows that psychotherapeutic treatment significantly reduce at one year follow-up dissociative experiences but also impulse regulation and body dissatisfaction, though alexithymia domains remained unchanged. People with EDs have a great deal of dissociative experiences^{6,14,39}. Intensive psychotherapeutic treatment may reduce dissociative proneness, supporting the patient to elaborating her vulnerability due to previous traumatic experiences establishing a link with them. Furthermore, giving freedom to speaking and listening seems producing a drop in impulsiveness, which can be communicated rather than acted. We value important also the decrease in body dissatisfaction, because it may play a role in EDs development and maintenance⁴⁰, but also may mediate on emotional factors, such as intolerance towards negative emotions⁴¹. In addition, two out the three AN-R cases showed an improvement in BMI. As regards alexithymia our findings could not show any change associated with psychotherapeutic treatment. Though alexithymia may well be a core psychological feature of EDS⁴², it has been already found it remains unchanged regardless of improvement in EDs symptoms⁴³, possibly because of its deep emotional nature.

We acknowledge several limitations of our study. First, given the exploratory nature of the study, the sample size was inevitably low, under powering our findings. In addition, scheduled one-year follow-up possibly struggles capturing long-term changes typical of psychotherapeutic treatment. Furthermore, though the sample is not homogeneous in terms of diagnostic features EDs, it may be seen with a “transdiagnostic model”^{44,45}, appropriate for a NHS, “real world” trial.

CONCLUSIONS

This work confirms the effectiveness of psychotherapy in people with EDs as regards dissociative moments, impulsivity and body dissatisfaction. On the contrary, alexithymia – a core psychological characteristic of the disorder – remains stable.

Table 1 - Descriptive characteristics of cases

Cases	Gender	Age	Axis I	Axis II	BMI Baseline	BMI 1 year follow-up	ED Previous Treatments
A	F	22	ANR	—	15.5	17.30	No
B	F	26	ANR	Obsessive	14.53	14.53	Yes
C	F	35	ANR	Histrionic	15.23	16.01	Yes
D	F	19	BN	Borderline	23.88	22.05	Yes
E	F	32	BN	—	22.86	21.57	Yes
F	F	22	NAS	Avoidant	25.39	21.87	No
G	F	45	NAS	Obsessive/ Schizoid	ND	ND	Yes
H	F	22	NAS	—	35.16	36.32	No

Legenda: ANR= Anorexia Nervosa Restricting Type; BN= Bulimia Nervosa; EDNOS= Eating Disorder Not Otherwise Specified; DP= Personality Disorder; BMI= Body Mass Index (kg/m2). BMI ≤17.5: significantly underweight; 17.6-18.9: underweight; 19.0-19.9: low weight; 20.0-24.9: normal weight; 25.0-29.9: overweight; ≥ 30: obesity. T1= Initial observation; T2= Final observation; ND= not determined because the patient refuse to weight.

Table 2. Core characteristics typical of EDs found in the whole clinical sample

Eating symptoms and food concerns about food, diet, eating
Body image disorder
Anxiety, depression, unhappiness
Emotional and impulse dysregulation or restricted emotional range
Unstable self-image
Fear of loneliness, of abandonment or rejection
Tendency to control or fluctuation between too much control or too little
Obsessiveness, concern for order and planning
Perfectionism
Feelings of emptiness
Shame
Feelings of inadequacy and inferiority, excessive self-criticism
Poor social skills and avoidance
Dependence
Dissociation proneness

Table 3. Pre- post-treatment differences at one year follow-up

	Baseline (T1)	One year follow-up (T2)	p T1 vs T2
	Mean (SD)	Mean (SD)	
EDI2	124.38 (60.25)	79.29 (64.43)	,18
DT-EDI	14.00 (7.13)	8.14 (7.84)	,09
BU-EDI	9.13 (9.51)	4.29 (7.67)	,11
BD-EDI	18.13 (8.03)	11.71 (9.01)	,02
IN-EDI	16.13 (9.34)	10.57 (10.34)	,18
P-EDI	4.13 (2.59)	4.00 (3.37)	,50
ID-EDI	7.25 (4.65)	7.29 (5.71)	,61
IA-EDI	11.13 (10.47)	5.43 (5.94)	,21
MF-EDI	10.00 (8.07)	9.29 (9.79)	,60
A-EDI	9.88 (5.41)	5.14 (2.97)	,08
I-EDI	13.00 (7.89)	4.86 (4.38)	,05
SI-EDI	11.63 (6.50)	8.57 (5.97)	,28
DES	22.38 (15.07)	6.53 (5.29)	,02
TAS20	60.08 (13.90)	53.00 (16.32)	,17
DIS	21.75 (10.02)	19.86 (5.61)	,87
DCS	15.38 (5.60)	14.86 (5.27)	,92
PO	19.50 (6.28)	18.29 (8.14)	,55

Further research, involving larger and homogenous samples, is needed in order to investigate the generalizability of these findings and to determine the role of dissociation, impulse regulation and alexithymia in EDs. The choice of meas-

uring instruments represents a relevant difficulty in methodological studies on alexithymia. As argued «the evaluation of the psychological trait that we want measure is inevitably dependent on the ability of the subject to report what he feels. Paradoxically, this ability is that should be insufficient in alexithymia»⁴⁶. It might be actually appropriate rethinking alexithymia in terms of different tools and modalities or respect the theoretical underlying conceptions⁴⁷.

Acknowledgements

Special thanks to Ombretta Fornasiero, Valentina Miot, Paola Pattini, the therapists who participated in the study and to Cristina Crocamo for statistical advice. Finally we thank the onlus foundation in Monza “Maria Bianca Corno Against Anorexia Nervosa”.

REFERENCES

1. Agras WS. The consequences and costs of eating disorders. *Psychiatr Clin North Am* 2001; 24: 371-9.
2. Hudson JI, Hiripi E, Pope HG, Kessler RC. The prevalence and correlates of eating disorders in the national comorbidity survey replication. *Biol Psychiatry* 2007; 61: 348-58.
3. Fernandez-Aranda F, Pinheiro AP, Thornton LM, et al. Impulse control disorders in women with eating disorders. *Psychiatry Res* 2008; 157: 147-57.
4. Becker AE, Grinspoon SK, Klibanski A, Herzog DB. Eating disorders. *N Engl J Med* 1999; 340, 1092-8.
5. Torem M. Dissociative states presenting as an eating disorder. *Am J Clin Hypn* 1986; 29: 137-42.
6. Demitrack MA, Putnam FW, Brewerton TD, Brandt HA, Gold PW. Relation of clinical variables to dissociative phenomena in eating disorders. *Am J Psychiatry* 1990; 147: 1184-8.
7. Putnam FW, Guroff JJ, Silberman EK, Barban L, Post RM. The clinical phenomenology of multiple personality disorder: review of 100 recent cases. *J Clin Psychiatry* 1986; 47: 285-93.
8. Chu JA, Dill LD. Dissociative symptoms in relation to childhood physical and sexual abuse. *Am J Psychiatry* 1990; 147: 887-92.
9. Tutkun H, Sar V, Yargic LI, Ozpulat T, Yanik M, Kiziltan E. Frequency of dissociative disorders among psychiatric inpatients in a Turkish university clinic. *Am J Psychiatry* 1988; 155: 800-5.
10. McCallum KE, Lock J, Kulla M, Rorty M, Wetzel RD. Dissociative symptoms and disorders in patients with eating disorders. *Dissociation* 1992; 5: 227-35.
11. Vanderlinden J, Van Dyck R, Vandereycken W, Vertommen H. Dissociative experiences and trauma in eating disorders. *Int J Eat Disord* 1993; 13: 187-93.
12. Everill J, Waller G, Macdonald W. Dissociation in bulimic and non-eating-disordered women. *Int J Eat Disord* 1995; 17: 127-34.
13. Abraham SF, Beumont PJV. How patients describe bulimia or binge eating. *Psychol Med* 1982; 12: 625-35.
14. La Mela C, Maglietta M, Castellini G, Amoroso L, Lucarelli S. Dissociation in eating disorders: relationship between dissociative experiences and binge-eating episodes. *Comprehensive Psychiatry* 2010; 51: 393-400.
15. Nagata T, Kiriike N, Iketani T, Kawarada Y, Tanaka H. History of childhood sexual or physical abuse in Japanese patients with eating disorders: relationship with dissociation and impulsive behaviours. *Psychol Med* 1999; 29: 935-42.
16. Vanderlinden J, Spinhoven P, Vandereycken W, van Dyck R. Dissociative and hypnotic experiences in eating disorder patients: an exploratory study. *Am J Clin Hypnoses* 1995; 38: 97-108.

Psychotherapeutic treatment of eating disorders may improve dissociative experiences and impulse regulation but not alexithymia

17. Paul T, Schroeter K, Dahme B, Nutzinger DO. Self-injurious behaviour in women with eating disorders. *Int J Eat Disord* 1995; 18: 99-102.
18. Taylor GJ, Bagby RM, Parker JDA. Disorders of affect regulation: alexithymia in medical and psychiatric illness. Cambridge: Cambridge University Press, 1997.
19. Bagby RM, Parker JDA, Taylor GJ. The twenty item Toronto Alexithymia Scale-1. Item selection and cross-validation of the factor structure. *J Psychosom Res* 1994; 38: 23-32.
20. Eizaguirre AE, De Cabezon AO, De Alda IO, Olariaga LJ, Maite J. Alexithymia and its relationship with anxiety and depression in eating disorders. *Pers Individ Differ* 2004; 36: 321-31.
21. Nicolò MA, Russo L. Una o più anoressie. Borla: Roma, 2010.
22. Markey MA, Vander Wal JS. The role of emotional intelligence and negative affect in bulimic symptomatology. *Compr Psychiatry* 2007; 48: 458-64.
23. Gratz KL, Roemer L. Multidimensional assessment of emotion regulation and dysregulation: development, factor structure, and initial validation of the Difficulties in Emotion Regulation Scale. *J Psychopathol Behav Assess* 2004; 26: 41-54.
24. Bulik CM, Klump KK, Thornton L, et al. Alcohol use disorder comorbidity in eating disorders: a multicentre study. *J Clin Psychiatry* 2004; 65: 1000-6.
25. National Collaborating Centre for Mental Health Eating Disorders. Core interventions in the treatment and management of anorexia nervosa, bulimia nervosa and related eating disorders. Leicester: The British Psychological Society, 2004.
26. Dalle Grave R, Calugi S, El Ghoch M, Conti M, Fairburn CG. Inpatient cognitive behavior therapy for adolescents with anorexia nervosa: immediate and longer-term effects. *Front Psychiatry* 2014; 12: 14.
27. Fairburn CG, Cooper Z, Shafran R. Cognitive behavior therapy for eating disorders: a transdiagnostic theory and treatment. *Behav Res Ther* 2003; 41: 509-28.
28. Priori M, Quagliata E. *Transfert e disturbi alimentari in età evolutiva*. Relazione presentata al Centro di Psicoanalisi Romano, 2005.
29. American Psychiatric Association. Diagnostic and statistical manual of mental disorders: DSM-IV-TR 4^a ed. American Psychiatric Publication, 2000.
30. Westen D, Shedler J, Lingardi V. La valutazione della personalità con la SWAP200. Milano: Raffaello Cortina Editore, 2003.
31. Garner DL, Olmstead MP, Polivy J. Development and validation of a multidimensional eating disorders inventory for anorexia nervosa and bulimia. *Int J Eat Disord* 1983; 2: 15-35.
32. Rizzardi M, Trombini E, Trombini G. EDI-2, Eating Disorder Inventory-2. Firenze: OS (Organizzazioni Speciali), 1995.
33. Bernstein EM, Putnam FW. Development, reliability, and validity of a dissociation scale. *J Nerv Ment Dis* 1986; 174: 727-35.
34. Steinberg M, Rounsaville B, Cicchetti D. Detection of dissociative disorders in psychiatric patients by a screening instrument and a structured diagnostic interview. *Am J Psychiatry* 1991; 148: 1050-4.
35. Taylor GJ, Bagby RM, Ryan D. Criterion validity of the Toronto Alexithymia Scale. *Psychosom Med* 1988; 50: 500-9.
36. Polivy J, Herman CP. Causes of eating disorders. *Ann Rev Psychol* 2002; 53: 187-213.
37. Parker JDA, Taylor GJ, Bagby RM. The Twenty Item Toronto Alexithymia Scale – III Reliability and factorial validity in a community population. *J Psychosom Res* 2003; 55: 269-75.
38. Cassin SE, von Ranson KM. Personality and eating disorders: a decade in review. *Clin Psychol Rev* 2005; 25: 895-916.
39. Iancu I, Cohen E, Yehudab YB, Kotlera M. Treatment of eating disorders improves eating symptoms but not alexithymia and dissociation proneness. *Compr Psychiatry* 2006; 47: 189-93.
40. Rohde P, Stice E, Marti CN. Development and predictive effects of eating disorder risk factors during adolescence: implications for prevention efforts. *Int J Eat Disord* 2015; 48: 187-98.
41. Koskina N, Giovazolias, T. The effect of attachment insecurity in the development of eating disturbances across gender: the role of body dissatisfaction. *J Psychol* 2010; 144: 449-71.
42. Guttman H, Laporte L. Alexithymia, empathy, and psychological symptoms in a family context. *Compr Psychiatry* 2002; 43: 448-55.
43. Schmidt U, Jiwanly A, Treasure J. A controlled study of alexithymia in eating disorders. *Compr Psychiatry* 1993; 34: 54-8.
44. Fairburn CG. *Cognitive behaviour therapy and eating disorder*. New York: The Guilford Press, 2008.
45. Dakanalis A, Carrà G, Calogero R, et al. Testing the cognitive-behavioural maintenance models across DSM-5 bulimic-type eating disorder diagnostic groups: a multi-centre study. *Eur Arch Psychiatry Clin Neurosci* 2014 Nov 22. [Epub ahead of print].
46. Porcelli P. Updates on the construct of alexithymia. 2005. www.psychomedia.it/pm/answer/psychosoma/porcelli-alex-2005.htm
47. Parling T, Mortazavi M, Ghaderi A. Alexithymia and emotional awareness in anorexia nervosa: time for a shift in the measurement of the concept? *Eat Behav* 2010; 11: 205-10.