

A Schema Therapy approach to complex dissociative disorder in a cross-cultural setting: a single case study

ALBERTO BARBIERI¹, FEDERICA VISCO-COMANDINI¹, ALESSANDRA TRIANNI¹, ANGELO MARIA SALIANI²

¹MEDU Psyché Center for Transcultural Mental Health, Rome, Italy; ²Scuola di Psicoterapia Cognitiva, Rome, Italy.

Summary. Complex dissociative disorders (CDD) include dissociative identity disorder (DID) and the most common other specified dissociative disorder (OSDD, type 1). While consensus-based treatments for CDD are lacking in several international guidelines, patients suffering from CDD show high levels of impairment, treatment utilization and costs. Migrants and refugees often present risk factors for trauma-related and dissociative disorders and need effective and culturally adapted treatments. Schema Therapy (ST) is an integrative psychotherapy that has been recently proposed as a treatment for CDD. This case study examined the process of individual ST, in a three phases-based approach, with a 38-year-old male Yemeni refugee with OSDD, type 1, posttraumatic stress disorder and borderline personality disorder. The treatment was provided in a Western country and the setting included two Western therapists (co-therapy) and an Arabic interpreter/cultural mediator. We assessed the patient's change by using self-report assessments of dissociative and PTSD symptoms, cognitive schemas over 2-year and 4-months treatment periods and a 6-months follow-up. Posttreatment and follow-up reliable change analyses showed significant improvements in dissociative and PTSD symptoms as well as in some cognitive schemas. Despite any firm conclusion cannot be drawn due to the limitations of this study (i.e., single case study), the findings suggest that ST integrated in a phase-oriented approach may be an effective treatment for CDD. Additionally, our study provides some preliminary elements about cross-cultural validity of the schema modes construct as well as cross-cultural effectiveness of ST. More research based on larger samples and specific cross-cultural focused design is needed to confirm these assumptions.

Key words. Case study, complex dissociative disorders, co-therapy, cross-cultural psychotherapy, schema therapy.

Un approccio di Schema Therapy al disturbo dissociativo complesso in un contesto interculturale: lo studio di un caso clinico.

Riassunto. I disturbi dissociativi complessi (CDD) includono il disturbo dissociativo dell'identità (DID) e il disturbo dissociativo con altra specificazione (OSDD, tipo 1). Mentre le terapie basate sul consenso per i CDD sono ancora assenti in diverse linee guida internazionali, i pazienti con CDD presentano oggi severi livelli di compromissione funzionale a fronte di trattamenti prolungati e costosi. Dal momento che le persone migranti e rifugiate presentano spesso fattori di rischio per i disturbi dissociativi e correlati a traumi, sono oggi necessari trattamenti efficaci e culturalmente adattati per questo gruppo di popolazione. La Schema Therapy (ST) è una psicoterapia integrata che è stata recentemente proposta come trattamento per i CDD. Questo studio presenta il caso clinico di una ST individuale, integrata in un approccio basato su tre fasi, di un rifugiato yemenita di 38 anni con OSDD, tipo 1, in comorbidità con un disturbo da stress post-traumatico e un disturbo borderline di personalità. Il trattamento ha avuto luogo in un paese occidentale con un setting comprendente due terapeuti occidentali (co-terapia) e un interprete/mediatore culturale arabo. Gli esiti della terapia sono stati periodicamente valutati attraverso misure self-report sia dei sintomi dissociativi e post-traumatici, sia degli schemi cognitivi nel corso di un trattamento di 2 anni e 4 mesi e di un follow-up di 6 mesi. Il reliable change index al post-trattamento e al follow-up ha mostrato miglioramenti significativi dei sintomi dissociativi e post-traumatici come anche di alcuni schemi cognitivi. Sebbene i limiti di questo studio (singolo caso clinico) non consentano di trarre conclusioni definitive, tuttavia i risultati suggeriscono che la ST possa essere un trattamento efficace per i CDD. Inoltre, questo studio fornisce alcuni elementi preliminari in favore della validità cross-culturale del costrutto degli schema modes e dell'efficacia interculturale della ST.

Parole chiave. Caso studio, co-terapia, disturbi dissociativi complessi, psicoterapia cross-culturale, schema therapy.

Introduction

According to the Diagnostic and Statistical Manual of Mental Disorders, fifth edition (DSM-5)¹, dissociative disorders are characterized by a disruption of the usually integrated functions of memory, identity, emotion, perception, body, representation, motor control, and behavior. Disruption of identity

includes confusion about one's identity as well as experiencing discrete and discordant senses of self, referred to as "identities", "personality states" or "dissociated self-states".

Complex dissociative disorders (CDD) include dissociative identity disorder (DID) and the most common dissociative disorder not otherwise specified (DDNOS, type 1), now known in the DSM-5 as

other specified dissociative disorder (OSDD, type 1)². The latter includes the subjects whose identity alteration is less marked to fully meet the diagnostic criterion of DID or who have identity alteration without dissociative amnesia. Patients suffering from CDD show high levels of impairment, high treatment utilization, and high treatment costs³. Numerous authors have demonstrated that pathological dissociation, dissociative disorders and possession phenomena occur at comparable rates in many different cultures or countries⁴. CDD in particular have been found in prevalence studies around the world using systematic assessments and validated interviews⁵, including countries of the Arabian Peninsula where the patient of this case study comes from⁶. One of the strongest predictors of dissociation is antecedent trauma, particularly early childhood trauma, as well as difficulties with attachment and parental emotional neglect⁷⁻¹⁰. The relationship between dissociation and many types of trauma is robust and has been validated across cultures in clinical and nonclinical samples by several studies⁷.

Emotion dysregulation is a core feature of CDD, as it is for borderline personality disorder (BPD). Indeed, there are many similarities between BPD and CDD, and dissociative disorders and BPD have often been reported to occur comorbidly¹¹⁻¹³. Childhood traumatic events (e.g., sexual abuse, physical and/or psychological abuse) and attachment difficulties which contribute to affect dysregulation are relevant risk factors for both borderline and dissociative symptoms^{9,14-17} along with biological diatheses or vulnerabilities. Posttraumatic stress disorder (PTSD) is also a common co-morbidity of patients with CDD¹⁸.

Currently, consensus-based treatments for CDD are lacking in several international guidelines (e.g., National Institute for Health and Care Excellence guidelines). Trauma and dissociation experts generally recommend an individual, phase-oriented psychotherapy using eclectically integrated interventions (e.g., psychodynamic, analytic, ego psychology, cognitive, behavioral, attachment, and neurodevelopmental approaches). As initially proposed by Pierre Janet in his pioneering work¹⁹, in the first phase, safety and symptom stabilization is established, in the second phase traumatic memories are confronted and processed, and in the third phase identity integration and rehabilitation is addressed (see practice-based guidelines by International Society for the Study of Trauma and Dissociation²⁰). Recently, some authors presented schema therapy (ST) as an alternative for phase-oriented trauma treatment models (in short 'POTT-models') for DID, considering POTT lengthy and with rather high reported dropout rates²¹. Although POTT length statistics are scarcely found, in their review Brand et al.¹⁸ reported a mean length of 8.4

(SD=4.8) years of treatment for CDD patients who were still in the last phase of treatment. ST^{22,23} is an integrative therapy for personality disorders that combines, in a systematic and structured model, elements of cognitive, behavioral, psychodynamic, and humanistic/experiential forms of psychotherapy. ST targets chronic emotional and cognitive maladaptive patterns, called "early maladaptive schemas" (EMS), which originate in adverse childhood experiences and early temperament. EMS are repeating themes about oneself and one's relationships that affect emotional processing, influence interpersonal style, and guide behavior. The primary objective of ST is to ameliorate EMS, replacing maladaptive coping responses with adaptive ones, and modify the transient, state-related manifestations of EMS, called "schema modes" which we refer to here as "modes". The latter are those EMS, coping responses or behaviors that are active at a given time for an individual²³ and are activated when specific EMS are triggered by specific situations, leading to overwhelming emotions and maladaptive coping responses that account for rapid changes in mood and behaviour often observed in personality disordered patients²⁴.

ST seems a viable option for the treatment of CDD given its emphasis on the consequences of early childhood neglect and abuse, and the explanation within the therapeutic model of the patient's experience of drastic shifts between personality states (i.e., mode switching or "flipping"). While a person is characterised by several schema modes, at a given moment in time, only one mode is predominant and determines the current behavior of patients with severe pathology, as this dominant mode shuts off the other modes²⁵. In this perspective, a dysfunctional mode can be considered as one of the aspects of the self that has not completely integrated with others in a cohesive personality structure and thus operates in a dissociated manner²⁶. Therefore, the various identities of a patient with CDD are regarded as extreme expressions of dissociated dysfunctional modes – especially in DID (in which a mode operate at a given moment without even being aware of the existence of the other modes) and in a milder way in OSDD, type 1. They differ from the modes of patients with personality disorders in how the patient experiences the mode, that is in the *degree* of experienced dissociation from the other modes^{23,25,26}. It has been proposed that the dissociation between different modes as described by Young et al.²³ is consistent with dissociation as conceptualized by the theory of structural dissociation of the personality (i.e., the trauma-related division of personality into different prototypical parts does not occur at random, but likely follows rather well-defined evolutionary prepared metaphorical "fault lines" in the structure of the personality)²⁷ and

concerns the concept of *désagrégation* (i.e., dis-integration) of the personality as originally advocated by Janet in his dissociation theory²⁸.

This study aims to contribute to the current literature by exploring the use of ST in the understanding and treatment of CDD in a cross-cultural setting consisting of a Yemeni refugee patient, an Arabic-speaking interpreter/cultural mediator and two Italian therapists. To the best of our knowledge there are no studies to date which investigated either the cross-cultural validity of schema modes or ST effectiveness in non-Western patients. We consider this case particularly relevant as the growing migratory flows in contemporary Europe increasingly pose the challenge of cross-cultural therapy settings and there is an urgent need to adopt integrated and culturally adapted psychotherapeutic approaches for migrants and refugees who often present trauma-related and dissociative disorders^{29,30}. We describe the 2-year and 4-months outpatient ST treatment and 6-months follow-up of a 38-year-old male Arabic-speaking refugee with CDD, PTSD and BPD. First, we will describe the treatment process. We will then present scores of the process measures ratings and the progress of the patient as repeatedly measured during treatment by the *Dissociative Experiences Scale - II* (DES-II)³¹, a self-report measure of EMS, and the Posttraumatic Stress Disorder Checklist (PCL-5)³². The rationale for applying ST to this specific case was that this is an evidence-based therapy³³ covering a wide array of consequences following adulthood and childhood trauma, including BPD^{34,35} and PTSD³⁶.

We adopted a therapeutic approach based on a transdiagnostic model, emphasizing common pathways of partly overlapping clinical syndromes such as CDD, PTSD and borderline phenomena^{21,37}.

Methods

PROCEDURE

Informed consent and psychotherapy process

The patient participated in the case study after giving written informed consent. He understood and agreed that we could make use of the therapists' notes, psychotherapy process, and measures for the purpose of the single case study. Additional informed consent was obtained for the audiotaped therapy sessions and for the publication of the drawings and paintings he donated to the therapists.

We used a fictitious name for the patient (i.e., Ali) and omitted all information that was not strictly necessary for understanding the case. The ST treatment process was documented on the basis of the therapists' case notes and extensive discussions between them.

PATIENT

Ali is a Yemeni 38-year-old painter and art teacher, married and childless. At the time of his first clinical interview, he was living in a reception center in Italy (a country of second resettlement for him after staying a few months in another European country following his escape from Yemen), waiting for the results of his asylum application. Ali reported that he had to flee his country due to persecution, while his wife remained in the country. In Yemen, the patient was a teacher at the Institute of Art in his city. As he used to portray naked men and women, he was threatened by some of the parents of his students who complained that his teaching methods were contrary to morality. Ali was detained for three days by the police following allegations of immoral behavior. During his detention, the patient was tortured and sexually abused by the police officers. After some years, he was attacked by another man on a motorcycle who shot him while he was driving his car.

As a child Ali was neglected and suffered from physical and psychological abuse by his mother who was totally dedicated to her daughter (Ali's stepsister), who suffered from psychosis. When Ali was 10 years old, his mother left the rest of the family to go abroad with her daughter where she remained for 13 years before returning. Ali reported that he was generally protected by his father, even though he was often violent with his children and also with him. The patient also suffered episodes of sexual abuse by other adults during his childhood. Ali was assessed using the Structured Clinical Interview for DSM-5 (SCID-5)³⁸ and the SCID-5 for Personality Disorders (SCID-5-PD)³⁹. The diagnosis of dissociative disorders was verified with the Dissociative Disorders Interview Schedule (DDIS)⁴⁰. The patient fulfilled DSM-5 criteria for OSDD, type 1 (i.e., chronic dissociative disorder with mixed symptoms), PTSD, BPD with narcissistic and histrionic traits.

SETTING AND PSYCHOTHERAPISTS

The case study took place at the Psyché Clinical Center for Transcultural Mental Health run by the medical humanitarian NGO Medici per i Diritti Umani (MEDU) in Rome (Italy). Dr. A.B., the case manager, and Dr. F.V.C., both certified cognitive-behavioral therapists, conducted the psychotherapy. Psychotherapy was given once a week in 60-minute sessions. All the first 17-months sessions were held in Arabic with the translation of an interpreter/certified cultural mediator native in Arabic which from now on we will refer to as the "interpreter". In the last stage, the sessions were held half in Italian and half in Arabic. The therapy sessions occurred with the two therapists present at the same time (i.e., co-therapy) and the interpreter. Indeed, psychotherapies supported by trained and experienced in-

terpreters are an effective option with traumatized refugees⁴¹. Some therapy sessions were audiotaped and examined during supervision with a certified ST supervisor and a certified cognitive-behavioral therapy (CBT) trainer with extensive ST experience (Dr. A.M.S.) providing feedback. Usually, trauma-focused treatments for PTSD in patients who have experienced multiple and/or prolonged interpersonal traumatic events (i.e., complex trauma) are made at the Psyché Clinical Center by two therapists. Co-therapy aims to prevent vicarious traumatization⁴² and compassion fatigue⁴³ through both mutual support and a diluted emotional drain for therapists who are faced with complex trauma cases that may be difficult to deal with alone⁴⁴. Both the therapists and the interpreter had a minimum of three years of experience in mental health work in cross-cultural settings.

MEASURES

The measures used in the study are displayed in the Supplemental table 1. A description of how using these standardized measures of process and outcome meets the American Psychological Association's criteria for evidence-based practice⁴⁵ is provided in the on-line supplemental materials. In line with the patient's symptoms and history as well as treatment adopted (i.e., ST approach), we used the translated Arabic versions of the Young Schema Questionnaire Short Form (YSQ-SF)⁴⁶, the PTSD Checklist for DSM-5 (PCL-5)⁴⁷, the Dissociative Experiences Scale-II (DES-II)⁴⁸. For any conceptual uncertainty on the items and/or on the Arabic terminology, the patient was supported by an independent interpreter (i.e., an interpreter different from the one who was part of the treatment team). For the therapy adherence ratings, we used the English version of the Schema Therapy Rating Scale (STRS)⁴⁹.

Table 1. Scores on the YSQ-SF Schema Domains, PCL-5 and DES-II Scores, Reliable Change Indices, and Effect Sizes for Pre-treatment to Posttreatment and Follow-Up.

Scales	Pretreatment	Midtreatment	Posttreatment	RCI	ES	Follow-up	RCI	ES
YSQ-SF Schema Domains								
Disconnection/rejection	3.6	2.0	0.2	3.79*	2.68	3.4	0.22	0.16
Impaired autonomy/performance	2.0	0.2	0.2	2.68*	2.34	1.2	1.19	1.04
Impaired limits	3.5	1.5	1.5	1.98*	1.68	3.0	0.50	0.42
Other-directedness	4.5	1.5	2.0	3.64*	3.00	2.0	3.64*	3.00
Over-vigilance/inhibition	3.5	2.5	2.0	1.97*	1.74	1.5	2.63*	2.32
PCL-5 total	42	23	22	3.62*	1.92	28	2.53*	1.34
Intrusion symptoms	8	1	2	2.79*	1.94	3	2.33*	1.61
Avoidance	5	1	1	3.52*	1.72	4	0.88	0.43
Negative alterations in cognition and moods	11	12	9	0.52	0.38	10	0.26	0.19
Alterations in Arousal	18	9	10	2.28*	1.74	11	1.99*	1.52
DES-II total	63.0	32.5	26.8	5.56*	2.84	18.2	6.82*	3.51
Taxon	70.0	35.0	12.5	3.14*	2.03	10.0	3.27*	2.12
Depersonalization/Derealization	53.3	18.3	10.0	1.99*	1.71	3.3	2.30*	1.98
Amnesic dissociation	65.0	20.0	28.7	2.25*	1.71	5.0	3.71*	2.83
Absorption & Imaginative Involvement	66.7	43.3	20.0	2.80*	2.69	36.7	1.80	1.73

Legend: RCI= Reliable Change Index; ES= Effect size; PCL-5= PTSD Checklist for DSM-5; DES-II= Dissociative Experiences Scale-II. *p<0.05.

Note: RCI scores for the schema domains, PCL-5 and DES-II have been reversed to correspond with positive treatment progress; higher scores on the PCL-5 and DES-II mean respectively more PTSD and dissociative symptoms, and higher scores on schema domains mean more maladaptive schemas.

Process measure ratings

To measure therapeutic alliance, we used the Helping Alliance Questionnaire (HAQ-II)⁵⁰. Since the Arabic version of this self-report questionnaire was not available, the independent interpreter translated each item out loud. The patient listened to each item and all the responses in Arabic and he then noted down his response. The interpreters attended a theoretical and practical training course on the use of all the study measures.

Results

PATIENT'S PRETREATMENT SCORES ON THE MEASURES

Ali obtained a pretreatment DES-II total score of 63.0. The DES-Taxon score (an eight-item measure drawn from the DES-II that determines whether the individual score is related to pathological dissociation) was 70.0. Table 1 provides the pretreatment scores for the YSQ-SF, PCL-5 total and sub-scales scores and the DES-II total and sub-scales scores.

PSYCHOTHERAPY PROCESS

Ali began psychotherapy six months after arriving in Italy and continued for two years and four months for a total of 85 sessions.

FIRST PHASE OF PSYCHOTHERAPY (STABILIZATION; FIRST FIVE MONTHS APPROXIMATELY)

Initial assessment

The stabilization phase lasted from session one to 17. During the first assessment sessions, Ali introduced himself as a very anxious person manifesting an apprehensive attitude evident from mimicry, gestures and language. Ali referred to "the loss of memory, getting lost, the boredom, the emptiness" as his main problems. He complained of frequent memory problems, difficulty in concentration and orientation, and rumination. In his opinion, "the loss of memory" started gradually since he was a child. The patient reported several recent episodes of amnesia concerning daily life events, especially in stressful situations, as well as impairment of episodic autobiographical memory (e.g., "I do not remember my marriage"). Moreover, the patient said that he often experienced an alteration in his sense of self (e.g., "During the day and during conversations I lose myself") and episodes of depersonalization (e.g., "I gave myself such strong slaps! It was as if another person had been slapped"). Ali also reported other three main problems. First, conversion symptoms, (i.e., temporary blindness episodes lasting about three minutes with concomitant strong

chest pains) and somatic symptoms including tingling, recurrent headaches and widespread pain, especially in the back, chest, legs and right side of his body. Second, he reported difficulties in interpersonal relationships. Ali was unable to establish or maintain significant bonds in the affective, social and work environment. In this regard, the patient reported having felt pervasive shame in social situations since his childhood as well as deep anger towards people who made him feel worthless. Third, he showed posttraumatic symptoms linked to the complex trauma experienced in adulthood, including intrusions, avoidances, arousal alterations. Ali declared to often think about suicide: "I hold back only because it is a sin for my Islamic religion".

After some months of therapy, Ali began to report frequent episodes in which internal dialogues between what he called "personalities" suddenly started in his mind. An example of how the patient described these internal dialogues is illustrated by the following: "A part of me says that I am a genius, another says that I am sick and that I am worthless. I have too many things inside, if I speak, I will explode. I am fighting, contradicting myself, I get so tired. Those who are inside me have been enemies for so long (ed., This last statement was accompanied by an emotional crisis with crying). These personalities of mine have been around for a long time. There is a conflict within me. There is a good person who loudly says: they believe in you! Then there is a deeper voice that says: they are humiliating you, they are exploiting you! Sometimes I enjoy watching these internal dialogues, sometimes they force me to confront myself as if I was in a meeting. I am like a father who has to hold off to my child personality, if I cannot hold off to that bad child, the child would say disgusting things. I have to prevail over him, I have to control him, but sometimes I listen to him". Ali mentioned sometimes the "bad child" as "the Devil".

The therapeutic alliance and the role of ancillary techniques

The stabilization phase sessions were mainly focused on improving patient's affect regulation and forming the therapeutic alliance as well as on ensuring that Psyché Center was perceived by Ali as a safe place. The therapists approached the patient with a welcoming, non-judgmental attitude, expressing empathy and supportive communication starting the limited reparenting process (see the following paragraph "The limited reparenting process" for details). Cognitive work and skills training were integrated in order to improve affect regulation and distress tolerance. The patient was trained to recognize the cognitive basis of his dysfunctional behavior and painful emotions, in particular through the execution of numerous exercises employing the ABC technique (i.e., a CBT technique to investigate

antecedents, cognitions, emotions and behaviors, of an event in the patient's life). The patient's emotions and beliefs were also investigated through the descriptions of the artistic works produced by the patient and presented by him during the sessions. The therapists instructed the patient to use some simple strategies for managing the most debilitating emotions such as anger and shame (e.g., relaxation techniques, memos on emotion awareness). The patient was trained in short daily grounding exercises to help him stay focused in the present reality.

Additionally, the therapists wrote a clinical report recognizing the patient as a survivor of torture. This report was important for the asylum application approval and helped to strengthen the patient's trust in the therapists. At the end of the stabilization phase, the therapeutic alliance was considerably strengthened (e.g., in session 12 Ali said: "At first I thought you were the secret service. Now I understand that you are serious"). The last sessions of this phase were focused on trauma and ST psychoeducation. The therapists trained the patient on the "language" of EMS and modes, and they worked in collaboration with him to develop a shared understanding of the patient's problems using these concepts.

SECOND PHASE OF PSYCHOTHERAPY (ELABORATION OF TRAUMATIC MEMORIES; 12 MONTHS APPROXIMATELY)

Retrieving childhood traumatic material

This phase lasted from session 18 to session 59. In session 18 the lifeline exercise was performed. The *lifeline* exercise is a tool developed by the Narrative Exposure Therapy (NET)⁵¹ as an effective first step towards discussing the traumatic material. The purpose of the lifeline is the reconstruction of subjectively significant life events in their chronological order. Notably, the patient showed a relative difficulty in retrieving specific events of his childhood as the lifeline appeared almost completely "empty" of positive or negative autobiographical events until his late adolescence.

The following sessions were dedicated to investigating Ali's childhood asking the patient to remember his relationship with both parents as well as events from his childhood biography. At this point of the therapy the patient started to say that he was subjected to frequent episodes of abuse and neglect by his mother: "My mother didn't touch me, didn't educate me, didn't show me any affection. She gave all the love and attention to my stepsister suffering from psychosis. My mother threatened me and beat me very hard. I was very afraid of her reactions". On the other hand, Ali tended to depict his father in a positive way, although he was sometimes described by the patient as a violent person. Ali began to use

writing and drawings as a homework to recall episodes of his childhood (e.g., sexual abuse episodes), bypassing the barrier of shame present in verbal communication. In subsequent sessions, the patient began to tell of a first sexual abuse that he suffered by an acquaintance of his father when he was nine (Supplemental figure 1-A1; in the drawing the patient re-enacts the events). The patient reported a following episode of sexual abuse by an adult when he was 15. Ali reported also consenting homosexual relationships during his adolescence, although he said he did not like that kind of relationship.

Case conceptualization according to ST

The case was conceptualized in terms of ST since the patient's "personalities" and their dissociative aspects could be framed in EMS and, subsequently, in poorly integrated modes. After some psychoeducation sessions on experiential techniques, Ali accepted to do imagery exercises which the therapists felt were essential for the change of EMS and to facilitate the integration of dissociated modes. Indeed, ST makes extensive use of experiential techniques, such as chair dialogues, guided imagery and imagery rescripting (IR)²³ in order to reprocess the patient's emotional distress originating from painful childhood experiences. In session 23, the therapists performed the imagery exercise of the safe place. In this exercise, the patient identified the Psyché Center room as *his* safe place. During session 24, a diagnostic imagery exercise was performed to identify specific relevant episodes of the past and present which could lead to the identification of the EMS. In the following session, the therapists examined with the patient his most relevant EMS (e.g., Abandonment, Defectiveness/Shame and Entitlement/Grandiosity) and coping strategies. Some sessions were focused on psychoeducation about coping modes (i.e., surrender, avoidant, overcompensator). The patient identified avoidance as his prevailing coping strategy. The patient was also trained to compile flash cards, that is, cards containing a small amount of information on EMS, as an aid to remember in daily life the most convincing arguments against EMS and the most adaptive behavioral strategies.

At this moment of the treatment, Ali donated to the therapists the painting named "Ali Child" (Supplemental figure 1-A2) where the patient represented himself as a little child (< three years old) clinging to the legs of the two therapists as if in search of care and protection from them. Then, the sessions were focused on talking about the modes of the Abandoned/Abused Child that Ali named "The Bud without Water" and the Detached Protector (i.e., avoidant coping mode that patient named "Ali the Solitary"). In session 35, the first IR was made on the episode of Ali rejected by his mother when he was

eight years old. In this memory, Ali sought the comfort of his mother but she abruptly pushed him away telling him that he should not cry and that she did not have time for him. During this exercise, Ali was able to introduce the Healthy Adult (formerly called the Counsellor and now named “Ali Kebir” which means adult in Arabic). In the following sessions, experiential exercises focused on the different Ali modes that gradually emerged (i.e., the Angry Child, the Abandoned/Abused Child, the Punitive Parent, the Detached Protector and the Overcompensator) were performed. The patient brought two more paintings. The first one depicted Ali Abandoned Child (Supplemental figure 1-A3; Ali represented himself as a 9 years old sad child waiting outside the closed door of the Psyché Center during the summer break). The second painting depicted “Ali the Solitary” (Supplemental figure 1-A4) represented as an adult in his room, isolated from the rest of the world. The painting also depicted aspects of the Overcompensator mode (“Ali the Genius”) showing a wall on which the patient painted the different ways he intended to commit suicide. In this case, suicide was seen by Ali as an extreme act to claim his own greatness against those who in Italy humiliated him by treating him badly.

After 10 months, the case conceptualization was complete. As Ali found the concept of modes easier to understand than EMS, the therapists shared with the patient a simplified version of the case conceptualization (figure 1) using only schema modes. Modes were grouped by their function, constantly focusing on the underlying needs. The right part of figure 1 shows the child and the punitive parent modes. Ali defined the Punitive Parent as “The Tough Mother”. The Abandoned/Abused Child often felt fear of abandonment and terror of abuse while at other times he rather felt shame, experiencing humiliation and unworthiness related to childhood experiences with his mother. The Angry Child was defined by Ali in the first sessions as “The Devil who says lousy things” and then as “The Volcano”. On the left side, the dysfunctional coping modes (i.e., the Detached Protector and the Overcompensator) are represented. At the beginning of phase 1, these modes were strong and rigid (particularly the Detached Protector) blocking access to the child and parent modes. In the upper left, a small circle represents the Healthy Adult at the beginning of ST. The therapists shared with the patient the therapeutic goal deriving from the modes’ conceptualization (i.e., a progressive integration of the dissociated modes towards a more cohesive sense of self) and the integration process of achieving it, that is, strengthening the Healthy Adult that in this way is able to give protection and love to the Abandoned/Abused Child, regulate the Angry Child, counter the Punitive Parent, and contain the dysfunctional coping modes.




Figure 1. Schema Mode Model used in Ali’s therapy.

Note. The size of the circles describing Ali’s modes are proportional to the relevance of each mode at the beginning of the therapy.

IR of the childhood abuse episode

In session 54, an IR was focused on the sexual abuse that Ali suffered from an acquaintance of his father when he was nine years old. During the exercise, Ali introduced the Angry Child who violently attacked the abuser with a series of punches against a pillow held by the interpreter. The Tough Mother then scolded the child. At this point, Ali was able to introduce the Healthy Adult (Ali Kebir) who partially consoled him. Afterwards, the therapist came into the room and fully consoled the Abused Child. The following is the central dialogue of the IR:

Dr. A.B.: What are you feeling at the moment Ali?

Ali: I am disgusted by this man. I am afraid, I am terrified. I am afraid he will kill me. I try to escape but he is strong and he’s holding me tight. I feel his lousy breath, his beard, his genitals. (Abused Child)

Dr. A.B.: What do you want right now?

Ali: I want help! May an adult help me, but there is no one around in this basement. I am afraid to scream because he will kill me! (Abused Child)

Dr. A.B.: At this moment the adult Ali enters the scene; Ali Kebir, what do you see?

Ali: I see a man abusing a child. I feel great anger. Even today when I see a strange adult approaching a child I remain alert. (Healthy Adult)

Dr. A.B.: What do you do and what do you say now?

Ali: I tell the child to run away and be safe at home (Healthy Adult). I do not tell that man anything, I cannot say anything. I beat him. I have so much anger that grows; I do not control it. I need to hit something around here for hours! (Angry Child) (The interpreter took a pillow and the patient repeatedly hit the pillow with a flurry of raging fists). (Angry Child)

Dr. A.B.: Now what are you saying to that man?

Ali: I want to insult him! (Ali begins to sob). That man is now on the ground, I still want to beat him; I want to destroy him. (Angry Child)

Dr. A.B.: Where is the child now? What do you do for the child?

Ali: He went away... I go to the father of the child and tell him to take care of him because he is a beautiful child and must be protected. (Healthy Adult)

Dr. A.B.: What else do you do?

Ali: I tell the child, "Why did you go with that man to the basement? Nobody will believe you! We do not behave like that; you have to pay attention!" (Punitive Parent)

Dr. A.B.: Do you want to say or do something else as an adult?

Ali: No

Dr. A.B.: How is Ali as a child now?

Ali: I feel safer, the fear is gone. But I feel lonely, I need an older brother, an adult who will not abandon me. (Abandoned Child)

Dr. A.B.: Can you do this as an adult for Ali as a child?

Ali: Yes, I caress the baby; I tell him "Do not be ashamed to tell your father things, but I will not always be there to help you. You have to trust other people too". (Healthy Adult)

Dr. A.B.: How does Ali child feel now?

Ali: A little better; but could you help me not to feel alone too? (Abandoned Child)

Dr. A.B.: Okay, I will enter the scene and I am close to you along with Ali Kebir. We are here to protect you and we will not leave you alone!

Ali: I feel good; I feel safer. I feel like I'm hugging both of you! (Happy Child)

In the following session, the patient stated that he had always felt a deep hatred for the abuser which diminished with the exercise. Afterwards, Ali reported of having been involved in numerous brawls from the ages of 20 to 28 because of his impulsiveness. Because of these fights he was detained by the police for some days for about seven times. Ali realized that thanks to therapy he had acquired greater self-control skills. Throughout the second year of therapy, Ali showed a progressive improvement in his relationships. The patient began to report daily life episodes in which he could better manage anger and shame by means of assertive strategies. Overall, Ali's social integration was proceeding positively. His knowledge of Italian was also improving. The patient reported a remarkable improvement in his dissociative and posttraumatic symptoms. In this phase, the IR on the sexual abuse episode was repeated; the patient introduced the Healthy Adult who was able to give protection and feel compassion and love towards the Abused Child without the need for the therapist to enter the scene.

The limited reparenting process

According to ST, the therapeutic relationship – conceptualized as *limited reparenting* – is a fundamental vehicle to bring about corrective emotional experiences^{23,52}. Through *limited reparenting*, the therapist creates an active, caring, parent-like relationship with the patient, while respecting the boundaries of the

therapeutic relationship. Although developing a limited reparenting relationship with a patient with CDD can be challenging, it is an essential component of the therapeutic process. Co-therapy with a male and female therapist increased the complexity of the therapeutic relationship. Following a situation in which Ali experienced unfairness that triggered his mistrust and abandonment schemas, an example of limited reparenting in co-therapy is provided here:

Ali: A few days ago, the director of my center went to my room when I was not there and found dirt because my roommates were never cleaning. I went to her to try to explain. I knocked on the door and she looked at me, she lowered her eyes and did not consider me. She told me: "Later, later, later!". I thought she did not want to talk to me. There was also a lawyer in her office. This lawyer never wanted to talk to me; I feel she does not want me. At that moment I was destroyed. I felt that there was no more space for me in that center. I felt that I was finished because before that moment I had given the director a little faith. I felt anger, anger! The heart was beating fast. Breathing became faster. My hands were shaking.

Dr. A.B.: I can imagine that you felt angry and mistreated, because you have been dismissed without the possibility of explaining. Did this episode remind you of similar moments in your childhood?

Ali: [He thinks for a short moment] Yes! I felt the same feelings when my mother first and my stepmother later did not want me and threw me out. My stepmother sometimes convinced my father that I had to stay away from him. This was unfair, it was terrible!

Dr. A.B.: I understand that these have been very painful moments for you as every child has the right to be welcomed and treated fairly by their parents. That is what you missed in your childhood.

Ali: Yes, you are right! The lawyer was happy that the director had kicked me out like when my father scolded me and my stepmother was happy for it.

Dr. A.B.: I understand how you must have felt. Isn't it possible, however, that the director was just very busy at the time, perhaps even under stress, and she just wanted to tell you to come back later?

Ali: Yes, but she could at least have greeted me and made me say a few words instead of throwing me in that disdainful way. I felt like human trash!

Dr. F.V.C.: I understand that at that moment you were asking for attention and to be welcomed. We want you to know that we are close to you and we want to help you to overcome these difficult situations and these painful emotions. We are by your side and genuinely interested in you as a person.

Since therapy required a long-term relationship with a patient who experienced very intense emotions and aroused equally strong ones in the people who were close to him, co-therapy was important in supporting therapists to manage their own EMS and modes, which were activated by the patient during the sessions. Further, co-therapy was helpful to acquire greater awareness of therapists' EMS and modes in peer-to-peer briefings between the sessions. The therapeutic relationship developed with different characteristics depending on the mode presented by the patient. Ali appeared initially distrustful, mainly

showing the avoidant coping mode (i.e., the Detached Protector). After the first months of treatment, Ali seemed fragile and in need of care. The Abandoned/Abused Child and the Punitive Parent were the most frequently present modes during the sessions. As the therapy progressed, Ali showed a growing trust in the therapists, and he also showed a tendency to idealise them. In the second phase, the patient progressively developed the ability to open up with the therapists, overcoming the avoidance of traumatic memories. The therapy gradually brought out new painful episodes of childhood that the patient urgently needed to communicate. In this phase, the Angry Child mode tended to emerge more easily, but also the Healthy Adult was progressively present with greater stability.

THIRD PHASE OF PSYCHOTHERAPY (INTEGRATION AND RESOCIALIZATION; ELEVEN MONTHS APPROXIMATELY)

This phase of the therapy was focused on consolidating therapeutic gains and in supporting Ali's progressive autonomy in his daily life. In order to encourage the autonomy of the patient towards the therapists, one of the two therapists (Dr. F.V.C.) began to progressively decrease her presence in the sessions. Over a period of about 10 months the sessions were reduced to one every two weeks. Throughout the third phase, Ali expanded progressively his social and professional network. Ali began to attend a carpentry internship and a cultural mediation course. He also did some job interviews. Relationship problems were less frequent in this phase. Most of the episodes concerned apparently minor situations (e.g., the queue at the dentist, the menu in the canteen, a reproach from the carpentry teacher) in which Ali felt unloved, alone and rejected. Such situations triggered the Angry Child's mode and emotions of helpless anger. Interventions such as role-playing and social skills training were used with the aim of promoting healthier relationships. Two-chair exercises focused on the dialogue between the Angry Child and the Healthy Adult were performed. The cognitive work was aimed at strengthening the Healthy Adult's perspective, always asking the following questions to the patient: "What would Ali Kebir say?" and "What would he do?". IR were performed on episodes of childhood domestic violence that emerged at this stage (e.g., the episode of Ali's brother beating his sister and their father wanting to kill them). At this stage, Ali reported a marked reduction of amnesia episodes in his daily life.

TERMINATION OF PSYCHOTHERAPY

Psychotherapy was interrupted in the third phase since Ali decided to move to northern Italy where he

had contacts with compatriots who could guarantee him a stable job. He wrote a long thank you message to greet the whole MEDU team. He also gave the therapists a painting that depicted him as a 10-year-old boy who, wearing a MEDU t-shirt, coming out happy from a pond where he played with other children: "I felt as if, thanks to therapy, I was happy as a child" (Supplemental figure 1-A5). Thus, the psychotherapy stopped when the process of traumatic memories retrieving as well as the change and the integration of the modes were still occurring.

FOLLOW-UP INTERVIEWS (SIX MONTHS AFTER THE END OF THERAPY)

Six months after ending ST, the therapists invited Ali for some follow-up interviews. Ali immediately agreed saying that he was happy to be able to speak again with the therapists who had helped him so much. After spending a few months in northern Italy, unhappy with the working conditions, he moved to another European country where he sought asylum. There, he lived in a refugee reception center. During the online interviews, he talked about his difficulties after ending therapy. When he was in this country, his mother died and he tested positive for covid-19. The patient described that difficult moment as follows: "I was desperate and jumped under a car. I did not get hurt but they took me to a psychiatric ward. I wanted to hurt myself a little and then come back alive. After this period of crisis, things are better now".

As requested by the therapists, Ali wrote the following evaluation in response to this question: how is it with your different sides (modes) at the moment? "Before answering this question, I would like to say something. Knowing how my personalities work gave me the motivation to understand the torment in which I lived and for which I had no explanation. This awareness has helped me a lot. To answer the question, I have noticed an overall improvement. Ali Kebir is now stronger and more present even if in the most difficult moments Ali the Abandoned Child still suffers. The positive changes due to therapy have been many and I'm not the only one to notice them, but also the people close to me. For example, I isolate myself much less from others".

QUANTITATIVE ASSESSMENT OF PSYCHOTHERAPY OUTCOME

In order to examine Ali's change from pre- to posttreatment and follow-up, we used the reliable change index (RCI)⁵². The RCI provides a z-score, where higher scores correspond with improvement and the threshold for significant improvement (at $p < .05$) lies at a z-score ≥ 1.96 . Table 1 provides the mid- and/or posttreatment and follow-up scores

on the YSQ-SF schema domains, PCL-5 total and subscale scores, DES-II total and subscale scores, the reliable change indices, and effect sizes. The description of the RCI calculation as well as the outcomes of each scale are provided in the supplemental materials.

PROCESS MEASURES RATINGS

The Helping Alliance (HAQ-II)

HAQ-II collects data from both the therapist and the patient to measure the strength of the bond between them⁴⁹. All the patient and therapists HAQ-II scores are displayed in table 2. The suggested cut-off point for good versus poor alliance was 4.57 mean score⁴⁹. While at the beginning of psychotherapy only one score out of four was above the cut-off (Therapist II versus Patient), in the second and third year all four scores were above the cut-off point indicating good therapeutic alliance between the patient and the therapists.

Discussion

This single case study is the first to describe a ST treatment for a CDD patient, with comorbid BPD and PTSD, in a cross-cultural setting. ST was used within a phase-oriented approach integrating CBT and NET techniques, skills training (e.g., grounding) as well as expressive modalities (i.e., artwork). Although it was interrupted in the third phase when the improvements were still ongoing, the therapy showed a partial effectiveness. The time to reach the last phase of treatment (2.4 years versus a mean of 8.4 years for other POTT-models)¹⁷ suggests a potentially shorter treatment duration compared to other phase-oriented approaches.

This case study is consistent with ST conceptualization of dissociation, in which CDD's different self-states can be conceptualized as extreme forms

of non-integrated dysfunctional modes. A schema mode can be defined according to the position it occupies along a dissociation spectrum from an extreme in which an individual is able to activate/express more than one mode at the same time (i.e., normal mood changes) to the opposite extreme in which an individual uses a mode, without having, at that moment, the awareness of the others (i.e., DID). Accordingly, the substantial difference between CDD identities and BPD modes would be the intensity and severity of the modes' dissociation. Taking into consideration this spectrum, Ali's complex dissociative disorder (i.e., OSDD, type 1) can be placed between BPD and DID, the severity and intensity of its modes' dissociation being higher than the former and lower than the latter. Recent studies have also shown that some typical modes of BPD present in Ali (e.g., the Angry Child, the Abandoned Child, the Detached Protector), would be able to significantly predict dissociative experiences^{53,25}. Furthermore, Johnston²⁵ showed how in the cases of BPD they studied it was not the childhood trauma *per se* that lead to the dissociative symptoms, but rather the dysfunctional modes originating from the childhood trauma. From a clinical and therapeutic point of view, the work of Johnston et al.²⁵ confirmed the importance of identifying and integrating the modes originating from the EMS (in particular the modes Angry Child and Abandoned Child) as well as strengthening the Healthy Adult in these patients. Accordingly, Ali's pervasive dissociative experiences would stem in dysfunctional modes (i.e., the different "personalities" recognized by Ali) originated in turn from neglect and childhood trauma. Moreover, the complex traumatic events occurred in Ali's adulthood would have acted as decompensation factors exacerbating the division of the parts of the personality (i.e., not-integrated dysfunctional schema modes) and consequently the dissociative symptoms.

In accordance with Brand et al.⁵⁴, we believe that ST for CDD should be integrated in a phase-oriented

Table 2. Helping Alliance Questionnaire Mean Scores (and SD) during the course of psychotherapy.

HAQ-II scores	Time during psychotherapy		
	First year	Second year	Third year
Patient			
Patient versus Therapist A.B.	2.73 (1.24)	5.18 (0.78)	5.31 (0.82)
Patient versus Therapist F.V.C.	3.00 (1.33)	5.15 (0.67)	5.26 (0.73)
Therapists			
Therapist A.B. versus Patient	4.05 (0.52)	5.15 (0.61)	5.21 (0.54)
Therapist F.V.C. versus Patient	4.68 (0.89)	5.22 (0.63)	5.31 (0.67)

approach (i.e., POTT model), and even more when, as it often occurs, BPD and PTSD occur comorbidly. In the case of Ali, creating a solid therapeutic alliance with both therapists (i.e., limited reparenting process) and improving patient affect regulation and coping skills were essential stabilization phase steps in order to apply the ST's mode approach as well as to face the elaboration of traumatic memories. Using a stabilization phase does not necessarily mean adopting an excessively long therapy or never reaching the stage of traumatic memories' elaboration as feared by Huntjens and colleagues⁵⁵. In the case of Ali, the stabilization phase lasted 17 sessions, just one more than the 16 sessions of preliminary education phase proposed in the Huntjens et al.'s⁵⁵ ST protocol for DID. Furthermore, the findings of this study highlight the importance of completing the third treatment phase (i.e., integration) when CDD, BPD and PTSD occur comorbidly. In the case of Ali, the significant improvements in all the EMS domains as well as in the prominent dissociative and posttraumatic symptoms reached at the termination of therapy were not fully maintained at the six-months follow-up. While the patient maintained significant improvements in two EMS domains, and in dissociative and posttraumatic symptoms, the improvements of some of the EMS domains were not. In particular, the Disconnection/Rejection domain score returned almost to pretreatment level. We assumed that not having completed the integration phase may subsequently have favored the re-emergence of the more rooted EMS (i.e., Disconnection/Rejection). Moreover, this case study confirms that the treatment of complex posttraumatic and dissociative disorders is often marked by recurrent relapses for which it is necessary to return from an advanced phase to a previous therapeutic stage (e.g., from integration phase to elaboration of traumatic memories) so that the therapy, rather than a linear progression, resembles a spiral path⁵⁶.

Another relevant element of the current case study refers to co-therapy which presents noteworthy aspects regarding the limited reparenting. Indeed, in the course of Ali's therapy, a double limited reparenting developed in which the two therapists (a woman and a man) took the role of the patient's parental figures (Supplemental figure 1-A2 "Ali Child"). This clinical case suggests that in a ST co-therapy, the two reparenting dyad (i.e., patient – male therapist and patient – female therapist) may act in a complementary and synergistic way allowing a solid therapeutic relationship, which is essential for all the three treatment phases of a CDD patient. In our case, the progressive and balanced growth of Ali's therapeutic alliance with both therapists is statistically highlighted by the HAQ-II administered in the first and third year of therapy: i.e. the scale scores of the patient versus the therapists

and vice versa were similar to each other presenting a specular evolution.

Dealing with the emotions related to traumatic experiences and countering the adult abuser through experiential work (i.e., IR) was a fundamental step of the therapy. In consideration of the patient's vulnerability and to prevent dissociative reactions, the technique IR was introduced gradually and with particular caution. The safe place and the imagery for assessment exercises were performed in the sixth month of therapy after the stabilization phase. The first IR was made after one year of therapy, while the episode of child abuse was dealt with much later during the second year of therapy, after having previously addressed it at a cognitive level.

To our knowledge, this is the first case study where ST was applied to the treatment of a CDD in a non-Western patient. The therapy was applied in a Western country (i.e., Italy) and the setting consisted of a Yemeni refugee patient, an Arabic-speaking interpreter/cultural mediator and two Italian therapists. Indeed, one of the most interesting aspects of our case study was related to the possible cross-cultural effectiveness of ST. In this regard, it is relevant to note that the patient had explicitly and spontaneously remarked several times, both during the therapy and the follow-up interviews, how well he recognized himself in the conceptualization of the case and in the approach of the modes, which he called "personalities". Although the role of the interpreter was essential in assigning the closest words to describe the emotions felt by the patient, we believe that the understanding of the characteristics and functions of the different modes were the result of a genuine elaboration of the patient. Of note, through his paintings and drawings, the patient was also able to visually represent some of his modes. Indeed, artwork was an important tool integrated to the treatment that facilitated the non-verbal communication of the modes, emotional states and traumatic experiences. The cultural adaptation of the therapy essentially consisted of language translation and considerations of cultural values and contextual stressors which can be considered more like cultural attunements⁵⁷. In this regard, a distinction must be made between adaptations in multicultural Western societies and non-Western societies. On the basis of the limited data of comparative experimental studies, adaptation/attunements efforts in communities that are in close contact with Western societies can be presumed more efficient⁵⁸. Therefore, even if a certain culturally adapted/attuned psychotherapy model is proved to be effective for a certain ethnic minority group in a Western country, this does not imply that this model is as effective in another country with populations of the same ethnic origin. Moreover, Pan, Huey, and Hernandez⁵⁹ notably showed that the difference between the effectiveness of culturally adapted psychotherapy and standard psycho-

therapy is less in more acculturated clients (as in the case of Ali), and vice versa. Therefore, the preliminary findings of this study on the cross-cultural effectiveness of ST should be considered with caution as their generalizability is limited by the features of this study (i.e., single case study), the characteristics of the setting (i.e., cross-cultural setting in a Western country) and the characteristics of the patient (i.e., Ali's specific educational and socio-cultural background).

The findings of our study have several limitations. First, it should be noted that the results of the measures administered to patients with CDD are often inconsistent, because these depend on which self-state participated in the assessment and on the fact that self-states may inhibit truthful responses⁶⁰. Future studies could use measures such as the integration measure (IM)⁶¹ to evaluate the state of personality integration during the course of therapy and at its termination. Second, very few studies with small samples investigated the validity and reliability of the Arabic versions of the DES-II and YSQ-SF measures. The scores on these questionnaires must therefore be interpreted with caution. Indeed, the cross-cultural adaptation of a psychological self-administered questionnaire is a complex task requiring that, the items must not only be translated appropriately linguistically, but also adapted culturally to maintain the content validity of the instrument at a conceptual level across different cultures⁶². Third, as there were no measures for the therapeutic alliance available in Arabic, we used the Italian version of the HAQ-II whose items were translated into Arabic by the interpreter for the patient when filling in the questionnaire. This process could have resulted in some minor differences in meaning and/or understanding during the translation as the full translation and back translation procedures to determine the quality of the scale for use in Arabic was not implemented. Furthermore, in our study, follow-up interviews took place six months after the termination of therapy. A longer-term follow-up would be necessary to confirm the stability of the changes achieved with the therapy.

Conclusions

Although any firm conclusion cannot be drawn as a consequence of various limitations which straiten our ability to generalize from these findings to other contexts, this case study suggests that ST integrated in a phase-oriented approach may be an effective treatment for CDD. Our study also provides some preliminary elements about cross-cultural validity of the schema modes construct as well as cross-cultural effectiveness of ST. This study therefore serves as a prime investigation of the effectiveness of ST approach to CDD in a cross-cultural setting in a Western country.

More research based on larger samples in different countries and/or specific design (e.g., studies designed to address cultural meanings of illness as well as cultural conceptualizations of treatment) is needed to confirm these assumptions. Of note, a case-series study to test the hypothesis that ST is an effective treatment for DID is currently in progress⁵⁵.

Disclosure: the data that support the findings of this study are available on request from the corresponding author.

Acknowledgments: we would like to express our gratitude to "Ali" for his consent to use his case for this study. Furthermore, we are grateful to Moez Chamki and Giulia De Lucia for their assistance in this study.

Conflict of interests: the authors have no conflict of interests to declare.

References

1. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th edition. Arlington, VA: American Psychiatric Publishing, 2013.
2. Brand BL, Lanius RA. Chronic complex dissociative disorders and borderline personality disorder: disorders of emotion dysregulation? *Borderline Personal Disord Emot Dysregul* 2014; 1: 13.
3. Myrick AC, Webermann AR, Langeland W, Putnam FW, Brand BL. Treatment of dissociative disorders and reported changes in inpatient and outpatient cost estimates. *Eur J Psychotraumatol* 2017; 8: 1375829.
4. Krüger C. Culture, trauma and dissociation: a broadening perspective for our field. *J Trauma Dissociation* 2020; 21: 1-13.
5. Sar V. Epidemiology of dissociative disorders: an overview. *Epidemiology Research International* 2011; 1-8.
6. Bener A, Saad AG, Micallef R, Ghuloum S, Sabri S. Sociodemographic and clinical characteristics of patients with dissociative disorders in an Arabian society. *Med Princ Pract* 2006; 15: 362-7.
7. Dalenberg CJ, Brand BL, Gleaves DH, et al. Evaluation of the evidence for the trauma and fantasy models of dissociation. *Psychol Bull* 2012; 138: 550-88.
8. Farina B, Liotti M, Imperatori C. The role of attachment trauma and disintegrative pathogenic processes in the traumatic-dissociative dimension. *Front Psychol* 2019; 10: 933.
9. Putnam KT, Harris WW, Putnam FW. Synergistic childhood adversities and complex adult psychopathology. *J Trauma Stress* 2013; 26: 435-42.
10. Schore AN. Attachment trauma and the developing right brain: origins of pathological dissociation. In: Dell PF, O'Neil JA (eds). *Dissociation and the dissociative disorders: DSM-V and beyond*. New York, NY: Routledge/Taylor & Francis Group, 2009.
11. Ross CA. Borderline personality disorder and dissociation. *J Trauma Dissociation* 2007; 8: 71-80.
12. Sar V, Akyuz G, Kugu N, Ozturk E, Ertem-Vehid H. Axis I dissociative disorder comorbidity in borderline personality disorder and reports of childhood trauma. *J Clin Psychiatry* 2006; 67: 1583-90.
13. Zittel Conklin C, Westen D. Borderline personality disorder in clinical practice. *Am J Psychiatry* 2005; 162: 867-75.
14. Herman JL. *Trauma and recovery: the aftermath of violence—from domestic abuse to political terror*. New York, NY: Basic Books, 1997.

15. Herman JL, Perry JC, van der Kolk BA. Childhood trauma in borderline personality disorder. *Am J Psychiatry* 1989; 146: 490-5.
16. Liotti G. Disorganized-disoriented attachment in the etiology of the dissociative disorders. *Dissociation: Progress in the Dissociative Disorders* 1992; 5: 196-204.
17. Van der Hart O, Nijenhuis ERS, Steele K. *The haunted self: structural dissociation and the treatment of chronic traumatization*. New York, NY: Norton, 2006.
18. Brand BL, Classen CC, Lanius R, et al. A naturalistic study of dissociative identity disorder and dissociative disorder not otherwise specified patients treated by community clinicians. *Psychological Trauma Theory Research Practice and Policy* 2009; 1: 153-71.
19. Janet P. *Névroses et idées fixes*. Paris: Félix Alcan, 1898.
20. International Society for the Study of Trauma and Dissociation. Guidelines for treating dissociative identity disorder in adults, third revision: summary version. *J Trauma Dissociation* 2011; 12: 188-212.
21. Huntjens R, Rijkeboer MM, Arntz A. Schema therapy for Dissociative Identity Disorder (DID): rationale and study protocol. *Eur J Psychotraumatol* 2019; 10: 1571377.
22. Young JE. *Cognitive therapy for personality disorders: a schema focused approach*. Sarasota, FL: Professional Resource Exchange, 1994.
23. Young JE, Klosko JS, Weishaar ME. *Schema therapy: a practitioner's guide*. London: Guilford Press, 2003.
24. Kellogg SH, Young JE. Schema therapy for borderline personality disorder. *J Clin Psychol* 2006; 62: 445-58.
25. Lobbstaal J, van Vreeswijk M, Arntz A. Shedding light on schema modes: a clarification of the mode concept and its current research status. *Neth J Psychol* 2007; 63: 76-85.
26. Johnston C, Dorahy MJ, Courtney D, Bayles T, O'Kane M. Dysfunctional schema modes, childhood trauma and dissociation in borderline personality disorder. *J Behav Ther Exp Psychiatry* 2009; 40: 248-55.
27. Steele K, van der Hart O, Nijenhuis ER. Phase-oriented treatment of structural dissociation in complex traumatization: overcoming trauma-related phobias. *J Trauma Dissociation* 2005; 6: 11-53.
28. Janet P. *The major symptoms of hysteria*. New York, NY: Macmillan. 1907
29. Palic S, Carlsson J, Armour C, Elklit A. Assessment of dissociation in Bosnian treatment-seeking refugees in Denmark. *Nordic J Psychiatry* 2015; 69: 307-14.
30. Van Ommeren M, de Jong JT, Sharma B, Komproe I, Thapa SB, Cardena E. Psychiatric disorders among tortured Bhutanese refugees in Nepal. *Arch Gen Psychiatry* 2001; 58: 475-82.
31. Carlson EB, Putnam FW. An update on the Dissociative Experiences Scale. *Dissociation* 1993; 6: 16-27.
32. Weathers F, Litz B, Keane T, Palmieri T, Marx BP, Schnurr P. The PTSD Checklist for DSM-5 (PCL-5). Scale available from the National Center for PTSD at: www.ptsd.va.gov [last accessed April 22, 2022].
33. Taylor C, Bee P, Haddock G. Does schema therapy change schemas and symptoms? A systematic review across mental health disorders. *Psychol Psychother* 2017; 90: 456-79.
34. Farrell JM, Shaw IA, Webber MA. A schema-focused approach to group psychotherapy for outpatients with borderline personality disorder: a randomized controlled trial. *J Behav Ther Exp Psychiatry* 2009; 40: 317-28.
35. Giesen-Bloo J, van Dyck R, Spinhoven P, et al. Outpatient psychotherapy for borderline personality disorder: randomized trial of schema-focused therapy vs transference-focused psychotherapy. *Arch Gen Psychiatry* 2006; 63: 649-58.
36. Cockram DM, Drummond PD, Lee CW. Role and treatment of early maladaptive schemas in Vietnam veterans with PTSD. *Clin Psychol Psychother* 2010; 17: 165-82.
37. Šar V. Parallel-distinct structures of internal world and external reality: disavowing and re-claiming the self-identity in the aftermath of trauma-generated dissociation. *Front Psychol* 2017; 8: 216.
38. First MB, Williams JWB, Karg RS, Spitzer RL. *Structured Clinical Interview for DSM-5® Disorders: Clinician Version (SCID-5-CV)*. Washington, DC: American Psychiatric Association Publishing, 2016.
39. First MB, Williams JWB, Karg RS, Spitzer RL. *Structured Clinical Interview for DSM-5® Personality Disorders (SCID-5-PD)*. Washington, DC: American Psychiatric Association Publishing, 2016.
40. Ross CA. *Multiple personality disorder: diagnosis, clinical features, and treatment*. New York, NY: Wiley, 1989.
41. Brune M, Eiroá-Orosa FJ, Fischer-Ortman J, Delijaj B, Haasen H. Intermediated communication by interpreters in psychotherapy with traumatized refugees. *Int J Cult Ment Health* 2011; 20: 144-51.
42. Sexton L. Vicarious traumatization of counsellors and effects on their workplaces. *Br J Guid and Couns* 1999; 27: 393-403.
43. Figley CR. Compassion fatigue as secondary traumatic stress disorder: an overview. In: Figley CR (ed). *Compassion fatigue: coping with secondary traumatic stress disorder in those who treat the traumatized*. New York, NY: Brunner/Mazel, 1995.
44. Russell A, Russell L. The uses and abuses of co-therapy. *J Marital Fam Ther* 1979; 5: 39-46.
45. American Psychological Association. Evidence-based practice in psychology. *American Psychologist* 2006; 61: 271-85.
46. Khater MEL, Gilany H, EL-Belsha M, Abdel-Moneim A. Reliability of the Arabic version of the Young Schema Questionnaire short-form among orphanage residents. *Middle East Journal of Psychiatry and Alzheimers* 2011; 2: 3-7.
47. Ibrahim H, Ertl V, Catani C, Ismail AA, Neuner F. The validity of Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5) as screening instrument with Kurdish and Arab displaced populations living in the Kurdistan region of Iraq. *BMC Psychiatry* 2018; 18: 259.
48. Al-Eithan M, Al Juban H, Robert AA. Dissociative experiences and their relationship to mood problems among Arab mothers of disabled children. *Pan Afr Med J* 2013; 15: 21.
49. Young JE, Fosse G. *Schema Therapy Rating Scale*. New York, NY: Cognitive Therapy Center of New York, 2005.
50. Luborsky L, Barber JP, Siqueland L, et al. The Revised Helping Alliance Questionnaire (HAQ-II): Psychometric Properties. *J Psychother Pract Res* 1996; 5: 260-71.
51. Schauer M, Neuner F, Elbert T. *Narrative Exposure Therapy: a short term treatment for Traumatic Stress Disorders (2nd edition)*. Cambridge, MA: Hogrefe Publishing, 2011.
52. Nordahl HM, Nysæter TE. Schema therapy for patients with borderline personality disorder: a single case series. *J Behav Ther Exp Psychiatry* 2005; 36: 254-64.
53. Jacobson NS, Truax P. Clinical significance: a statistical approach to defining meaningful change in psychotherapy research. *J Consult Clin Psychol* 1991; 59: 12-9.
54. Barazandeh H, Kissane DW, Saeedi N, Gordon M. Schema modes and dissociation in borderline personality disorder/traits in adolescents or young adults. *Psychiatry Res* 2018; 261: 1-6.
55. Brand BL, Loewenstein RJ, Schielke HJ, et al. Cautions and concerns about Huntjens et al.'s Schema Therapy for Dissociative Identity Disorder. *Eur J Psychotraumatol* 2019; 10: 1631698.

56. Huntjens R, Rijkeboer MM, Arntz A. Schema therapy for Dissociative Identity Disorder (DID): rationale and study protocol. *Eur J Psychotraumatol* 2019; 10: 1571377.
57. Chu JA. *Rebuilding shattered lives: the responsible treatment of complex post-traumatic and dissociative disorders*. New York, NY: Wiley, 1998.
58. Falicov CJ. Commentary: on the wisdom and challenges of culturally attuned treatments for Latinos. *Fam Process* 2009; 48: 292-309.
59. Koç V, Kafa G. Cross-cultural research on psychotherapy: the need for a change. *J Cross Cult Psychol* 2019; 50: 100-15.
60. Pan D, Huey SJ, Hernandez D. Culturally adapted versus standard exposure treatment for phobic Asian Americans: treatment efficacy, moderators, and predictors. *Cultur Divers Ethnic Minor Psychol* 2011; 17: 11-22.
61. Van der Hart O, Nijenhuis ERS, Steele K. *The haunted self: structural dissociation and the treatment of chronic traumatization*. New York, NY: Norton, 2006.
62. Barlow MR, Chu JA. Measuring fragmentation in dissociative identity disorder: the integration measure and relationship to switching and time in therapy. *Eur J Psychotraumatol* 2014; 5.
63. Beaton DE, Bombardier C, Guillemin F, Ferraz MB. Guidelines for the process of cross-cultural adaptation of self-report measures. *Spine* 2000; 25: 3186-91.

Supplemental materials

Introduction

CROSS-CULTURAL ISSUES RELATED WITH ALI'S THERAPY

Mental health in Yemen has developed within a context of wars, internal struggles, poverty, high birth rates and illiteracy. Mental health disorders are often connected to myth, superstition and religious beliefs¹. Nevertheless, mental health in Yemen has benefited from government support, even if modest, and human resource development projects allowing the development of several mental health public services and private clinics mainly used by urban population. According to a survey study, paranoia with schizophrenia was the most common diagnosis, followed by emotional disturbances (e.g., depression and anxiety)². Suicide is still a highly sensitive issue in Yemen, and stigma and shame are connected to it¹. As per WHO estimates from 2018, suicide mortality rate is 8.5 per 100,000 people³.

To date, few evidence-based psychological treatments have been tested for this culture, although some institutional and individual initiatives have developed intervention models that demonstrate the effectiveness of mental health services in the Yemeni context¹. The literacy rate among the population aged 15 years and older (54.1%)⁴ reflects a socio-culturally

heterogeneous society with a marked difference between large urban centers and rural areas. The patient in our study had a high level of education and, although he had always lived in his country until his escape to Europe, he had a good knowledge of the Western culture as he taught Western Renaissance art in his city's art school. In this regard, the patient himself spontaneously requested a professional psychological support once he arrived in Italy without this being a stigmatizing factor for him.

Method

MEASURES

With the aim to meet the American Psychological Association's criteria for Evidence-Based Practice⁵ as well as the *Clinical Utility* dimension in the Criteria for Evaluating Treatment Guidelines⁶, we integrated verbatim clinical case material with standardized measures of process and outcome evaluated at different times across treatment. Consequently, this case study includes the assessment of three standardized outcome measures (YSQ-SF for changes in EMS) and target symptom (DES-II for dissociative symptoms and PCL-5 for PTSD symptoms), as well as one process measure (HAQ-II for therapeutic alliance) evaluated on at least three separate occasions. Specific outcome data were presented using standardized mean difference (i.e., effect size) and clinical significance methodology (i.e., reliable change index)⁷.

Supplemental Table 1. List of measures used in the case study, when administered, and by whom.

Topic	Measures	Description	Time in treatment	Rated by
Dissociative symptoms	Dissociative Experiences Scale-II (DES-II) ⁹	The DES-II is a 28-item self-report measure that assesses dissociative symptoms	Pre-, mid-, posttreatment and follow-up	Self-report administered with an independent interpreter support
Early maladaptive schemas	Young Schema Questionnaire Short Form (YSQ-SF) ⁸	The YSQ-SF is a self-report assessment instrument of 75 items, to assess the 15 EMS	Pre-, mid-, posttreatment, and follow-up	Self-report administered with an independent interpreter support
Post-traumatic Stress Disorder	PTSD Checklist for DSM-5 (PCL-5) ⁹	The PCL-5 is a 20-item self-report measure that assesses the 20 DSM-5 symptoms of PTSD	Pre-, mid-, posttreatment, and follow-up	Self-report administered with an independent interpreter support
Therapeutic alliance	Helping Alliance Questionnaire (HAQ-II) ¹²	The HAQ-II is a 19 item self-report questionnaire that measures the strength of the client therapist alliance	Pre-, mid-, and posttreatment	Self-report administered with an independent interpreter support
Therapy adherence	Schema Therapy Rating Scale (STRS) ¹³	The STRS measures the psychotherapist competency in using ST	Two audiotaped sessions during the second year of psychotherapy	An independent rater, a certified ST therapist

Note. Pretreatment= at the start of treatment (first phase); midtreatment =beginning of second year of treatment (second phase); posttreatment= beginning of third year of treatment, except for the YSQ-SF, which was completed after 2-year and 4-months of treatment (third phase); follow-up= 6 months after ending treatment.



Supplemental Figure 1. Drawings and paintings made by Ali relating to therapy: A1) Re-enactment of the sexual abuse at the age of 9; A2) Ali Child with the therapists; A3) Ali Abandoned Child; A4) Ali the Solitary; A5) Ali Happy Child.

Results

QUANTITATIVE ASSESSMENT OF PSYCHOTHERAPY OUTCOME

To calculate the RCI, we used reliability coefficients and standard deviations from validation studies for YSQ-SF8 and PCL-59. For the DES-II, we used the reliability coefficients and the standard deviations obtained in a sample of 10 Arabic-speaking asylum seekers who sought treatment at the Psyché Clinical Center for trauma-related psychological distress. A small study of reliability of the scale was carried out on the 10 asylum seekers. Cronbach's α coefficients calculated for the DES-II scale showed high alpha value ($\alpha=.87$). Cronbach's α values for DES-II sub-scales (Taxon, Amnesic Dissociation, Depersonalization/Derealization, Absorption & Imaginative Involvement) were $\alpha=.79$, $\alpha=.71$, $\alpha=.63$, and $\alpha=.54$, respectively. Effect sizes (Cohen's d) were calculated as the difference between pretreatment and posttreatment or follow-up means divided by the standard deviations for the YSQ-SF, PCL-5, and DES-II from normative samples (as mentioned above).

Early maladaptive schemas (YSQ-SF). The scores on the YSQ-SF decreased significantly from pre- to posttreatment for all the EMS domains. From pre-treatment to follow-up, the decrease in the score re-

mained significant for two of the five EMS domains: Over-vigilance/Inhibition and Other-directedness.

Post-traumatic Stress Disorder (PCL-5). PCL-5 total score decreased significantly from pre- to posttreatment and follow-up to the point that at posttreatment and follow-up, Ali no longer met the DSM-5 criteria for the diagnosis of PTSD and the total scores (respectively 22 and 28) were below the PCL-5 indicative cut-off (33) for PTSD¹⁰.

Dissociative symptoms (DES-II). The total score of the DES-II and each of its subscales decreased significantly from pre- to posttreatment. At post-treatment, the total score (26.8) was below the indicative cut-off (30) for a dissociative disorder¹¹. The decrease in score also remained significant at follow-up. Notably, the total and each subscale scores recorded a further decrease at follow-up with the exception of the Absorption & Imaginative Involvement subscale.

References

1. Qasem Saleh MAB, Makki AM. Mental health in Yemen: obstacles and challenges. *Int Psychiatry* 2008; 5: 90-2.
2. YMHA (Yemeni Mental Health Association) [Survey] Al-Seha-AlAqilia 2006, No. 29-30.
3. WHO (2021, April 9). World Health Statistics data visualisation dashboard. Retrieved from <http://apps.who.int/gho/data/>

4. UNESCO (2021, April 9). Yemen. Retrieved from <http://uis.unesco.org/en/country/ye>
5. APA Presidential Task Force on Evidence-Based Practice. Evidence-based practice in psychology. *Am Psychol* 2006; 61: 271-85.
6. American Psychological Association. Criteria for evaluating treatment guidelines. *Am Psychol* 2002; 57: 1052-9.
7. Jacobson NS, Roberts LJ, Berns SB, McGlinchey JB. Methods for defining and determining the clinical significance of treatment effects: description, application, and alternatives. *J Consult Clin Psychol* 1999; 67: 300-7.
8. Khater MEL, Gilany H, EL-Belsha M, Abdel-Moneim A. Reliability of the Arabic version of the Young Schema Questionnaire short-form among orphanage residents. *Middle East Journal of Psychiatry and Alzheimers* 2011; 2: 3-7.
9. Ibrahim H, Ertl V, Catani C, Ismail AA, Neuner F. The validity of Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5) as screening instrument with Kurdish and Arab displaced populations living in the Kurdistan region of Iraq. *BMC Psychiatry* 2018; 18: 259.
10. Weathers F, Litz B, Keane T, Palmieri T, Marx BP, Schnurr P (2013). The PTSD Checklist for DSM-5 (PCL-5). Scale available from the National Center for PTSD at www.ptsd.va.gov
11. Carlson EB, Putnam FW, Ross CA, et al. Validity of the Dissociative Experiences Scale in screening for multiple personality disorder: a multicenter study. *Am J Psychiatry* 1993; 150: 1030-6.
12. Luborsky L, Barber JP, Siqueland L, et al. The Revised Helping Alliance Questionnaire (HAQ-II): psychometric properties. *J Psychother Pract Res* 1996; 5: 260-71.
13. Young JE, Fosse G. Schema Therapy Rating Scale. New York, NY: Cognitive Therapy Center of New York, 2005.