A preliminary study on similarities and dissimilarities of Neurological Soft Signs in schizophrenic and obsessive-compulsive disorders suggests a common maldevelopmental model

GIUSEPPE BERSANI¹, ADELE QUARTINI^{2,3}, GIORGIANA MANUALI⁴, FRANCESCA PACITTI⁵, ANGELA IANNITELLI⁵⁻⁷

¹Minds in Network; ²Department of Medical-Surgical Sciences and Biotechnologies, Faculty of Pharmacy and Medicine, Sapienza University of Rome; ³DSM ASL/LT, Unit of Psychiatry A. Fiorini Hospital, Terracina (LT); ⁴UOC SPDC Pertini Hospital, ASL RM2; ⁵Department of Clinical Sciences and Applied Biotechnology, University of L'Aquila, Italy; ⁶Psychoanalytical Society (SPI), Rome, Italy; ⁷International Psychoanalytical Association (IPA), London, UK.

Summary. Aim. Neurological Soft Signs (NSS) represent minor neurological signs related to non-specific cerebral alterations. They have been documented in many psychiatric disorders including schizophrenia (SCZ) and obsessive-compulsive disorder (OCD). Aim of this study was to determine and compare the incidence and severity of NSS in patients with SCZ, in patients with OCD, and healthy control subjects (HCs). Methods. Using the Neurological Evaluation Scale (NES), this study investigated NSS in 15 SCZ patients, 14 OCD patients, and 15 HCs. PANSS and Y-BOCS were used to evaluate clinical picture in both groups. Results. Patients with SCZ showed significantly higher scores compared to HCs in the NES total and each of the three NES subscales (Integrative Sensory Function, Motor Coordination, and Sequencing of Complex Motor Acts). Patients with OCD also showed significantly higher scores compared to HCs in the NES total, Motor Coordination and Sequencing of Complex Motor Acts, but not in Integrative Sensory Function. No significant differences emerged in the NES total and the various subscales scores between the two patients' groups. Conclusions. Our results seem to confirm the presence of NSS in both SCZ and OCD. The different types of NSS presented by the two patients' groups versus HCs further supports the findings of widespread cerebral alterations in SCZ, on the other hand, with a preferential involvement of prefrontal and frontal cortex in OCD.

Key words. Cerebral maldevelopment, neurodevelopment, neurological soft signs, obsessive-compulsive disorder, schizophrenia.

Introduction

Neurological Soft Signs (NSS) are subtle but observable impairments in motor and sensory functions that are not restricted to a characteristic of any Lo studio preliminare sulle similitudini e differenze dei Neurological Soft Signs suggerisce un comune modello di alterazione dello sviluppo nei disturbi schizofrenici e ossessivo-compulsivi.

Riassunto. Scopo. I Neurological Soft Signs (NSS) rappresentano dei segni neurologici cosiddetti minori correlati ad alterazioni cerebrali non specifiche. I NSS sono presenti in molti disturbi psichiatrici inclusi la schizofrenia (SCZ) e il disturbo ossessivo-compulsivo (OCD). Obiettivo dello studio è stato quello di determinare e comparare l'incidenza e la gravità dei NSS in pazienti con schizofrenia e in pazienti con OCD confrontandoli con controlli sani. Metodi. I NSS sono stati valutati in 15 pazienti schizofrenici, 14 pazienti con OCD e 15 controlli sani, attraverso la somministrazione della Neurological Evaluation Scale (NSS). La sintomatologia clinica è stata misurata attraverso la somministrazione della PANSS e della Y-BOCS. Risultati. I pazienti con schizofrenia hanno valori significativamente più alti nella NES totale e in tre sottoscale (Integrative Sensory Function, Motor Coordination e Sequencing of Complex Motor Acts) rispetto ai controlli sani. Anche i pazienti con OCD hanno, rispetto ai controlli sani, valori più alti nella NES totale e in due sottoscale (Motor Coordination e Sequencing of Complex Motor Acts) ma non nella Integrative Sensory Function. Nessuna significativa differenza è emersa tra i due gruppi di pazienti nel valore totale e nelle sottoscale della NES. Conclusioni. I risultati del nostro studio confermano la presenza di NSS in entrambi i gruppi di pazienti. Inoltre, i differenti tipi di NSS presenti nei due gruppi di pazienti studiati, rispetto ai controlli sani, supportano il dato di diffuse alterazioni cerebrali nei pazienti con schizofrenia e di un coinvolgimento preferenziale della corteccia frontale e prefrontale nei pazienti con OCD.

Parole chiave. Alterato sviluppo cerebrale, disturbo ossessivo-compulsivo, neurological soft signs, neurosviluppo, schizofrenia.

specific neurological condition¹ and typically, they are classified into signs relating to sensorimotor integration, motor coordination, sequencing of complex motor tasks, and disinhibition².

NSS have been proposed to be a marker of abnor-

mal brain development^{3,4}, and are known to correlate with a wide range of neurocognitive and neuroanatomical abnormalities, and it has been proposed that they represent an underlying defect in neural integration⁵.

NSS have been reported in excess in SCZ, where they are considered a neuro-dysfunction associated to this disorder⁶, with frequency ranging from 50% to 65% in contrast to 5% in normal population^{7,8} and they have been described both in first-episode drugnaïve and treated patients9-11. They have been associated with negative or deficit symptoms, cognitive functioning, an earlier onset of the illness and a poorer long-term outcome^{5,7,12}. The neural substrates of NSS in SCZ remain poorly understood. Basal ganglia and cerebellum, as somatomotor and somatosensory regions, would seem to be involved in impairment in visual processing and spatial orientation, as well as dysfunction of frontoparietal and cerebellar networks would seem to be implicated in the pathophysiology of NSS. It remains unclear if white matter integrity deficits or neurometabolic alterations contribute to NSS in the illness. Furthermore, only little is known about the temporal evolution of NSS, and in antipsychotic medication-naïve patients These data are very important gaps in the knowledge of NSS13.

On the whole, the etiology of NSS is uncertain; Tsuang et al.¹⁴ have argued that they might reflect genetic and non-genetic processes underpinning the predisposition to psychotic illness. Corroborating this assumption, NSS have well been documented in subjects genetically at risk of developing SCZ¹⁵. Indeed, according to the "Neurodevelopmental model", SCZ is believed to be the result of an interaction between genetic and environmental factors. More specifically, this model hypothesizes that, in subjects with SCZ, cerebral damage - caused by non-genetic factors, such as obstetric complications – eventually interacts with genetic liability to cause neurodevelopmental abnormalities; thus, the onset of SCZ in adult life might ensue when the brain matures sufficiently to call into operation the damaged system¹⁶.

With regard to OCD, controversial reports are present in the literature about the relationship between OCD and NSS. Some Authors have found a higher prevalence of NSS in patients with OCD¹⁷⁻¹⁹ and cognitive deficits in the domains of visuospatial ability, executive function, attention, and working memory in these patients compared to controls²⁰⁻²² yet other studies have failed to detect such differences²³⁻²⁵. Moreover, whilst some Authors^{17,25} suggest NSS as indicators of non-structural brain damage, which may cause OCD symptoms¹⁷, other Authors believe that NSS are likely to be related to both structural and functional brain abnormalities, especially regarding left basal ganglia and insula, and a disorder-specific fronto-striato-insular dysregulation in the form of poor frontal control over overactive basal ganglia²⁶, and that the affected regions of the brain for NSS probably resemble those in OCD²⁷. Whether NSS can be consider a marker of a deviant neurodevelopment in patients with OCD, as for patients with SCZ, is still a matter of debate²⁵.

Considering the former, the aim of the present pilot study was to determine the incidence and severity of NSS in patients with SCZ as compared with patients with OCD and a control group, and to evaluate differences in the expression of NSS in SCZ compared with OCD.

Materials and methods

Three groups of subjects took part in this study: 1) 15 patients with chronic SCZ, 2) 14 patients with OCD, and 3) 15 HCs. Psychiatric diagnoses were determined according to DSM-V criteria. Demographic information and past and current symptom history of all patients were obtained in semi-structured interviews. Clinical status was assessed by using the Scale for the Assessment of Positive and Negative Syndrome Scale (PANSS)²⁸ and the Yale Brown Obsessive Compulsive Scale (Y-BOCS)²⁹. Subjects were matched for age, sex and other demographic variables. Informed consent was obtained from all participants. In order to evaluate a possible correlation between NSS and previous and/or ongoing antipsychotic medications, a careful investigation was carried out with regard to the antipsychotic treatment throughout the illness course. Patients were recruited from the Unit of Psychiatry A. Fiorini Hospital in Terracina (Sapienza University, Rome). Inclusion criteria for all subjects were: 1) 15-60 years of age (in order to avoid bias due to degenerative diseases); 2) no previous or ongoing substance abuse; and 3) no previous or ongoing neurological disorders (such as intellectual disability, dementia, and/or history of traumatic brain injury with loss of consciousness).

NSS were assessed using an Italian version of the Neurological Evaluation Scale (NES)². This is a twenty-six items check list, fourteen of which are bilateral, and consists of three subscales relatively to the three different NSS cluster categories and their most likely, putative neuroanatomical localization⁷. Each item, except for the snout reflex and the suck reflex (evaluated with a score of 0 or 2), has a score scaled three points: 0= no anomaly; 1= slight presence of anomaly; 2= marked presence of anomaly. The NES has been employed following the instructions provided by the Authors. We took into consideration the score obtained at the NES total, as an index of global NSS seriousness, and the scores obtained at the three NES subscales: Integrative Sensory Function (the score of which is the sum of the items tactile extinction on bilateral stimulation, graphestesia, stereognosis, right/ left confusion and audio-visual integration); Motor Coordination (the score of which is the sum of the items tandem walk, finger-to-nose test, finger-thumb opposition and dysdiachokinesis); and Sequencing Of Complex Motor Acts (the score of which is the sum of the items fist-ring test, fist-edge-palm test and Ozeretski's test).

STATISTICAL ANALYSES

Analyses were carried out using SPSS V17 (SPSS Inc., Chicago, IL, USA). Sociodemographic clinical variables were described and compared amongst the three groups (OCD, SCZ, and HCs) with t tests, one-way ANOVAs or chi-square tests as appropriate. Pearson's correlations between the NES, the PANSS, and the Y-BOCS total and subscales scores were computed after testing data for normality and linearity distributions.

Because of the possible influence of antipsychotic treatment on NSS and because there was a significant difference in the age ratio, ANCOVAs, with age and lifetime antipsychotic treatment (duration of illness by the chlorpromazine equivalent units of current daily antipsychotic dose) as covariates, were performed for the NES total and subscales scores to evaluate contrasts amongst the groups. When data could not be transformed to meet the assumptions of normality, Kruskal-Wallis tests or Mann-Whitney tests were used as appropriate. The critical level for statistical significance was set at p<0.05. The exploratory nature of the study and the small sample size, which might have easily lead to a Type II error, argued against the application of correction for multiple comparisons.

Results

Demographic and clinical characteristics of the study participants are shown in table 1. SCZ patients were significantly older than the OCD patients and the HCs [F(2,42)=4.297, p=.020. Posthoc test: SCZ vs OCD: p=.020; SCZ vs HC: p=.012]. Age at onset was significantly earlier [t(27)=2.636], p=.014] in the OCD group than in the SCZ group. In the SCZ group, all patients were currently on antipsychotics. In the OCD group, 69.23% of patients (N=9) were currently on antidepressants and 30.77% (N=4) were currently being treated with antipsychotics. There were no significant differences for the remaining descriptive variables amongst the three subjects' groups [sex: $\chi^2(2)$ =.968, p=.616; educational level: F(2,42)=.908, p=.412; and duration of illness: t(27)=-.593, p=.558].

In the SCZ patients, the NES total score was positively correlated with the PANSS General Psychopathology Symptoms score (r=.808, p=.028). A positive correlation was also found between the right-sided tests score of the Integrative Sensory Function subscale of the NES and the PANSS-Negative symptoms score (r=.839, p=.018) and even between the rightsided tests score of the Integrative Sensory Function subscale of the NES and the PANSS total score (r=.952, p=.028). In the OCD patients, the NES total

Table 1. Characteristics of patients and control subjects.											
	SCZ (N=15)		OCD (N=14)		HCs (N=15)						
	Mean	SD	Mean	SD	Mean	SD	р				
Age (years)	41.47	12.48	32.14	9.37	31.36	8.72	0.02				
Educational level (years)	11.47	2.80	12.57	3.06	12.93	3.32	0.41				
Onset	38.93	15.58	28.93	8.50			0.01				
Duration of illness (months)	31.33	25.10	37.71	32.561	-	-	0.56				
Y-BOCS - Obsessions - Compulsions - Total	- - -	- -	13.36 9.36 22.73	2.38 3.58 5.31	- -	- - -	-				
PANSS - Positive - Negative - Total	17.00 27.71 92.71	5.164 5.678 15.861	- - -	- - -	- - -	- - -	-				
Antipsychotic treatment	15		9		0		-				

Legend: SCZ= patients with schizophrenia; OCD= patients with obsessive-compulsive disorders; HCs= healthy controls; Y-BOCS= Yale-Brown Obsessive Compulsive-Scale (subscores and total scores); PANSS= Positive And Negative Syndrome Scale (subscores and total scores).

and subscales scores did not correlate significantly with the Y-BOCS total or subtotal scores for obsessions or compulsions.

The NES total and subscales scores were significantly different amongst the three groups [NES total: F(2,42)=6.693, p=.003; Integrative Sensory Function: H(2)=15.514, p<.001; Motor Coordination: H(2)=12.945, p=.002; and Sequencing Of Complex Motor Acts: H(2)=13.326, p=.001] (table 2). Post-hoc tests revealed that the NES total score was significantly higher in both SCZ and OCD patients' groups than in the HCs group (respectively, p=.002 and p=.008).

With regard to SCZ patients, they performed higher than HCs at each of the three NES subscales scores (Integrative Sensory Function: U=19.500, z=-3.858, p<.001; Motor Coordination: U=34.000, z=-3.405, p=.001; and Sequencing Of Complex Motor Acts: U=21.000, z=-3.780, p<.001) and also at each of their right- (Integrative Sensory Function: U=56.000, z=-2.857, p=.004; Motor Coordination: U=56.000, z=-2.857, p=.004; and Sequencing Of Complex Motor Acts: U=42.000, z=-3.360, p=.001) and left-sided tests scores (Integrative Sensory Function: U=56.000, z=-2.861, p=.004; Motor Coordination: U=56.000, z=-2.851, p=.004; and Sequencing Of Complex Motor Acts: U=46.000, z=-2.890, p=.004). Apart from the Integrative Sensory Function subscale score, OCD patients performed higher than HCs at the various NES subscales scores (Motor Coordination: U=59.000, z=-2.170, p=.030 and Sequencing ff Complex Motor Acts: U=52.000, z=-2.267, p=.023). Interestingly, significant differences were found in both the right- and left- tests of the Sequencing of Complex Motor Acts

(U=56.000, z=-2.695, p=.007 and U=53.500, z=-2.403, p=.016, respectively), whereas, with regard to the Motor Coordination, this difference was seen only for the tests assessing the left side (U=56.000, z=-2.696, p=.007).

OCD and SCZ patients did not differ significantly in total scores on the NES or in any of the individual tests.

Discussion and conclusions

The first major finding of this study is the presence of NSS in both SCZ and OCD patients, although with significant differences in the type of NSS presented by the two patients' groups when compared to HCs.

According to previous studies⁵, the performances of SCZ patients were significantly worse than HCs in the NES total and each of the three NES subscales (Integrative Sensory Function, Motor Coordination, and Sequencing of Complex Motor Acts). Sequencing of Complex Motor Acts reflects the functions of the prefrontal cortex⁷; Motor Coordination signs are reported to be hypothetically related to fronto-cerebellar impairments^{7,30} and Integrative Sensory Function reflects the functions of the parietal lobe^{7,30}.

With regard to OCD patients, they also showed significantly higher scores compared to HCs in the NES total, Motor Coordination and Sequencing of Complex Motor Acts, but not in Integrative Sensory Function.

Such results seem to suggest widespread cerebral alterations in SCZ, on the other hand, with a preferential involvement of prefrontal and frontal cortex in

sive disorders and in control subjects.												
	SCZ (I	V=15)	OCD (N=14)		HCs (N=15)		р					
Domain	Mean	SD	Mean	SD	Mean	SD	SCZ vs. HCs	OCD vs HCs				
Sensory Integration - right-sided tests - left-sided tests - total score	0.73 0.93 3.67	0.88 1.10 2.74	0.50 0.93 1.71	0.76 1.07 1.90	0.00 0.00 0.43	0.00 0.00 0.85	0.004 0.004 <0.001	n.s. n.s. n.s.				
Motor Coordination (fronto-cerebellar) - right-sided tests - left-sided tests - total score	0.73 1.00 2.33	0.88 1.31 2.13	0.21 0.57 0.93	0.43 0.76 1.14	0.00 0.00 0.14	0.00 0.00 0.36	0.004 0.004 0.001	n.s. 0.007 0.030				
Motor sequencing (prefrontal) - right-sided tests - left-sided tests - total score	1.40 1.53 4.47	1.55 1.46 3.78	0.86 1.14 3.29	1.10 1.23 3.50	0.00 0.21 0.50	0.00 0.58 0.85	0.001 0.004 <0.001	0.007 0.016 0.023				
Total NES score	16.60	11.10	9.64	6.18	2.43	2.03	0.002	0.008				

Table 2. Scores on the Neurological Evaluation Scale (NES) in patients with schizophrenia, in patients with obsessive-compulsive disorders and in control subjects.

Legend: SCZ= patients with schizophrenia; OCD= patients with obsessive-compulsive disorders; HCs= healthy controls; SD= standard deviation; n.s.: not significant. OCD. As regards to laterality and cerebral dominance, our research did not identify an unequivocal laterality pattern of neurological signs in SCZ. Indeed, patients with SCZ performed worse than HCs in both the right- and left-sided tests. With regard to OCD patients, they also showed significant differences in both the right- and left-sided tests of Sequencing of Complex Motor Acts, whereas in Motor Coordination this difference was significant only for tests assessing the left side.

Some studies have reported for SCZ a predominance of neurological abnormalities on the right side of the body, but there are conflicting reports of higher scores for NSS on the left side of the body³¹.

Regarding OCD, few studies have investigated the laterality of NSS in these patients. Confirming our results, a trend for higher scores of the NSS on the left side, especially motor signs, was reported³².

The second major finding of our study is the lack of any significant differences in the NES total and/ or the various subscales scores between the two patients' groups. SCZ and OCD are disorders involving the prefrontal-striato-thalamic pathway, even though the cortical areas implicated in these illnesses are different: in SCZ, the main cortical area involved is the dorsolateral prefrontal cortex, in OCD it is the orbitofrontal cortex³³. The NES scale does not specifically assess orbitofrontal cortex functions and this could explain the lack of significant differences between the two groups.

As would be expected, the two patients' groups were different with respect to psychotropic administration. ANCOVAs, with age and lifetime antipsychotic treatment (duration of illness by the chlorpromazine equivalent units of current daily antipsychotic dose) as covariates did show an absolute independence of NSS from antipsychotic treatment. This result seems to confirm other studies that found no influence of medication on NSS in either SCZ or OCD^{9,11,17}.

NSS are considered signs of impaired neurodevelopment^{14,19,25}, thus the lack of any significant differences in the NES total and/or the various subscales scores between the two patients' groups seems to support a neurodevelopmental origin not only for SCZ, but also for OCD. It can be assumed that these neurological signs may depend on some functional brain abnormalities, potentially involved as substrate for SCZ and/or OCD. Genetic and non-genetic factors may cause brain alterations responsible for slight neurological abnormalities, then – in high-risk patients – they may precipitate a psychotic or an OCD symptomatology^{14,16,19}.

To conclude, both SCZ and OCD are associated with an increased rate of NSS, though with significant differences in the type of NSS presented by the two patients' groups when compared to HCs. The lack of any significant differences in the NES total and/or the various subscales scores between the two patients' groups may support a common neurodevelopmental origin.

The main limit of this study is the small sample size and the impossibility to completely exclude a potential confounding effect of psychotropic medication in patients with SCZ. Yet, our study confirms previous research further suggesting a possible extension of the "Neurodevelopmental Model" to SCZ, OCD and other mental diseases.

Conflict of interests: the authors have no conflict of interests to declare.

References

- 1. Rigucci S, Dimitri-Valente G, Mandarelli G, et al. Neurological soft signs discriminate schizophrenia from bipolar disorder. J Psychiatr Pract 2014; 20: 147-53.
- Buchanan RW, Heinrichs DW. The neurological evaluation scale (NES): a structured instrument for the assessment of neurological signs in schizophrenia. Psychiatry Res 1989; 27: 335-50.
- Bourgou Gaha S, Halayem Dhouib S, Amado I, Bouden A. Signes neurologiques mineurs dans la schizophrénie précoce. Encephale 2015;41: 209-14.
- Gay O, Plaze M, Oppenheim C, et al. Cortex morphology in first-episode psychosis patients with neurological soft signs. Schizophr Bull 2013; 39: 820-9.
- Chan RC, Xu^T, Heinrichs RW, Yu Y, Wang Y. Neurological soft signs in schizophrenia: a meta-analysis. Schizophr Bull 2009; 36: 1089-104.
- Mohr F, Hubmann W, Cohen R, et al. Neurological soft signs in schizophrenia: assessment and correlates. Eur Arch Psychiatry Clin Neurosci 1996; 246: 240-8.
- 7. Bombin I, Arango C, Buchanan RW. Significance and meaning of neurological signs in schizophrenia: two decades later. Schizophr Bull 2005; 31: 962-77.
- 8. Heinrichs DW, Buchanan RW. Significance and meaning of neurological signs in schizophrenia. Am J Psychiatry 1988; 145: 11-8.
- 9. Gupta S, Andreasen NC, Arndt S, et al. Neurological soft signs in neuroleptic-naïve and neuroleptic treated schizophrenic patients and in normal comparison subjects. Am J Psychiatry 1995; 152: 191-6.
- Dazzan P, Lloyd T, Morgan KD. Neurological abnormalities and cognitive ability in first-episode psychosis. Br J Psychiatry 2008; 193: 197-202.
- 11. Bersani G, Gherardelli S, Clemente R. Neurologic soft signs in schizophrenic patients treated with conventional and atypical antipsychotics. J Clin Psychopharmacol 2005; 25: 372-5.
- Bersani G, Clemente R, Gherardelli S, Pancheri P. Deficit of executive functions in schizophrenia: relationship to neurological soft signs and psychopathology. Psychopathology 2004; 37: 118-23.
- 13. Samson GD, Lahti AC, Kraguljac NV. The neural substrates of neurological soft signs in schizophrenia: a systematic review. Schizophrenia (Heidelb) 2022; 8: 42.
- 14. Tsuang MT, Faraone SV. The concept of target features in schizophrenia research. Acta Psychiatr Scand Suppl 1999; 395: 2-11.
- Marcus J, Hans SL, Lewow L, Wilkinson L, Burack CM. Neurological findings in high risk children: childhood assessment and 5-year follow-up. Schizophr Bull 1985; 11: 85-100.

- 16. Bersani G, Quartini A, Manuali G, et al. Influence of obstetric complication severity on brain morphology in schizophrenia: an MR study. Neuroradiology 2009; 51: 363-71.
- 17. Hollander E, Schiffman E, Cohen B, et al. Signs of central nervous system dysfunction in obsessive-compulsive disorder. Arch Gen Psychiatry 1990; 47: 27-32.
- Bolton D, Gibb W, Lees A, et al. Neurological soft signs in obsessive compulsive disorder: standardised assessment and comparison with schizophrenia. Behav Neurol 1998; 11: 197-204.
- 19. Rosenberg DR, Keshavan MS. A.E. Bennett Research Award. Toward a neurodevelopmental model of obsessive-compulsive disorder. Biol Psychiatry 1998; 43: 623-40.
- 20. Dhuri CV, Parkar SR. Soft neurological signs and cognitive function in obsessive-compulsive disorder patients. Indian J Psychol Med 2016; 38: 291-5.
- Jaafari N, Fernández de la Cruz L, Grau M, et al. Neurological soft signs in obsessive-compulsive disorder: two empirical studies and meta-analysis. Psychol Med 2013; 43: 1069-79.
- 22. Mergl R, Hegerl U. Neurological soft signs in patients with obsessive-compulsive disorder Fortschr Neurol Psychiatr 2005; 73: 504-16.
- Stein DJ, Hollander E, Simeon D, Cohen L, Islam MN, Aronowitz B. Neurological soft signs in female trichotillomania patients, obsessive-compulsive disorder patients, and healthy control subjects. J Neuropsychiatry Clin Neurosci 1994; 6: 184-7.
- Poyurovsky M, Faragian S, Pashinian A, et al. Neurological soft signs in schizophrenia patients with obsessivecompulsive disorder. J Neuropsychiatry Clin Neurosci 2007; 19: 145-50.

- 25. Jaafari N, Baup N, Bourdel MC, et al. Neurological soft signs in OCD patients with early age at onset, versus patients with schizophrenia and healthy subjects. J Neuropsychiatry Clin Neurosci 2011; 23: 409-16.
- Carlisi CO, Norman LJ, Lukito SS, et al. Comparative multimodal meta-analysis of structural and functional brain abnormalities in autism spectrum disorder and obsessive-compulsive disorder. Biol Psychiatry 2017; 82: 83-102.
- 27. Guz H, Aygun D. Neurological soft signs in obsessivecompulsive disorder. Neurol India 2004; 52: 72-5.
- Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS) for schizophrenia. Schizophr Bull 1987; 13: 261-76.
- Goodman WK, Price LH, Rasmussen SA, et al. The Yale-Brown Obsessive-Compulsive Scale. I. Development, use, and reliability. Arch Gen Psychiatry 1989; 46: 1006-11.
- Bersani G, Paolemili M, Quartini A, et al. Neurological soft signs and cerebral measurements investigated by means of MRI in schizophrenic patients. Neurosci Lett 2007; 413: 82-7.
- Dazzan P, Murray RM. Neurological soft signs in firstepisode psychosis: a systematic review. Br J Psychiatry 2002; 181: 50-7.
- Thomas N, Tharyan P. Soft neurological signs in drug-free people with schizophrenia with and without obsessivecompulsive symptoms. J Neuropsychiatry Clin Neurosci 2011; 23: 68-73.
- 33. Venkatasubramanian G, Rao NP, Behere RV. Neuroanatomical, neurochemical, and neurodevelopmental basis of obsessive-compulsive symptoms in schizophrenia. Indian J Psychol Med 2009; 31: 3-10.

Corresponding author: Dr. Francesca Pacitti E-mail: francesca.pacitti@univaq.it