

Diagnostic value and correlation analysis of deregulated oxy-Hb disorder in cognitive impairment in patients with alcohol dependence

JINGLIN BAI¹, XIUQIN HUANG¹, ZHANPING MA¹, XIANGYANG ZHENG¹, BOWEN JI¹

¹Department of Psychiatry, Hengshui Seventh People's Hospital, Hengshui, China.

Summary. Aim. To explore the expression of oxygenated hemoglobin (oxy-Hb) in the brain of patients with alcohol dependence. Further, to evaluate the diagnostic value of oxy-Hb in the cognitive impairment of patients with alcohol dependence. **Methods.** Seventy-one alcohol-dependent adults and 70 healthy adults were recruited in the observation and control groups, respectively. The oxy-Hb level and scores of Wisconsin card sorting test, brief visuospatial memory test-revised, digital sequence test, digital span test, and continuous performance test were recorded and compared between the two groups. Further, the sensitivity and specificity of the oxy-Hb level for the diagnosis of cognitive impairment in patients with alcohol dependence were evaluated by drawing an ROC curve. Moreover, the correlation between the oxy-Hb level and cognitive function test scores was analyzed by Pearson's correlation analysis. **Results.** During cognitive activities, the oxy-Hb levels in the observation group were lower than those in the control group ($p < 0.001$). The area under the curve (AUC) of oxy-Hb was 0.930, and the sensitivity and specificity were 89.1% and 84.0%, respectively. The scores of responses administered, trials to compete first category, perseverative responses, and the perseverative response errors in executive function in the observation group were higher than those in the control group ($p < 0.001$), whereas the correct responses score and categories completed score were lower than those in the control group ($p < 0.001$). Similarly, the memory and attention scores of the observation group were lower than those of the control group ($p < 0.001$). Moreover, the scores of correct responses, categories completed, brief visuospatial memory test-revised, digital sequence test, digital span test, and continuous performance test were positively correlated with the level of oxy-Hb ($p < 0.05$). However, the scores of responses administered, trials to compete first category, and perseverative response errors were negatively correlated with the oxy-Hb level ($p < 0.001$). **Conclusions.** The oxy-Hb levels are reduced in patients with alcohol dependence and have high diagnostic accuracy for cognitive impairment in such patients.

Key words. Alcohol-dependent, cognitive impairment, diagnosis, oxy-Hb.

Valore diagnostico e analisi di correlazione del disturbo ossi-Hb deregolato nel deterioramento cognitivo nei pazienti con dipendenza da alcol.

Riassunto. Scopo. Esplorare l'espressione dell'emoglobina ossigenata (ossi-Hb) nel cervello di pazienti con dipendenza da alcol. Inoltre, valutare il valore diagnostico di ossi-Hb nel deterioramento cognitivo dei pazienti con dipendenza da alcol. **Metodi.** Settantuno adulti alcol-dipendenti e 70 adulti sani sono stati reclutati rispettivamente nei gruppi di osservazione e di controllo. Il livello di ossi-Hb e i punteggi del test di selezione di carte di Wisconsin, del test di memoria visuospatiale breve rivisto, del test di sequenza digitale, del test di span digitale e del test delle prestazioni continue sono stati registrati e confrontati tra i due gruppi. Inoltre, la sensibilità e la specificità del livello di ossi-Hb per la diagnosi di deterioramento cognitivo nei pazienti con dipendenza da alcol sono state valutate disegnando una curva ROC. La correlazione tra il livello di ossi-Hb e i punteggi dei test delle funzioni cognitive è stata analizzata dall'analisi di correlazione di Pearson. **Risultati.** Durante le attività cognitive, i livelli di ossi-Hb nel gruppo di osservazione erano inferiori a quelli del gruppo di controllo ($p < 0,001$). L'area sotto la curva (AUC) dell'ossi-Hb era 0,930 e la sensibilità e la specificità erano rispettivamente dell'89,1% e dell'84%. I punteggi delle risposte somministrate, le prove per competere nella prima categoria, le risposte perseverative e gli errori di risposta perseverativa nella funzione esecutiva nel gruppo di osservazione erano superiori a quelli del gruppo di controllo ($p < 0,001$), mentre il punteggio delle risposte corrette e il punteggio delle categorie completate erano inferiori a quelli del gruppo di controllo ($p < 0,001$). Allo stesso modo, i punteggi di memoria e attenzione del gruppo di osservazione erano inferiori a quelli del gruppo di controllo ($p < 0,001$). Inoltre, i punteggi delle risposte corrette, delle categorie completate, del breve test di memoria visuospatiale rivisto, del test di sequenza digitale, del test di span digitale e del test continuo delle prestazioni sono stati positivamente correlati con il livello di ossi-Hb ($p < 0,05$). Tuttavia, i punteggi delle risposte somministrate, le prove per competere nella prima categoria e gli errori di risposta perseverativa erano correlati negativamente con il livello di ossi-Hb ($p < 0,001$). **Conclusioni.** I livelli di ossi-Hb sono ridotti nei pazienti con dipendenza da alcol e hanno un'elevata accuratezza diagnostica per il deterioramento cognitivo in tali pazienti.

Parole chiave. Deterioramento cognitivo, diagnosi, dipendenza da alcol, ossi-Hb.

Introduction

In the current social scenario, most individuals do not have a correct understanding of alcohol consumption, which results in alcohol abuse and eventually leads to alcohol dependence. Alcohol dependence is a physiological and psychological state induced by repeated alcohol consumption, manifested by a strong craving for alcohol and forced experience of frequent alcohol consumption^{1,2}. Patients with alcohol dependence find it challenging to effectively control the occurrence, beginning, end, and dosage of alcohol consumption behavior³. After complete or partial abstinence, alcohol withdrawal symptoms, such as tremors, transient hallucination, seizures, and delirium, often appear, and these symptoms rapidly disappear after recover from alcohol consumption⁴. According to a study published in *The Lancet Medical Journal*, alcohol consumption has increased by 10% from 1990 to 2017, and it is considered that if this trend continues, alcohol consumption will increase by 17% over the next 10 years⁵. Alcoholism can seriously endanger human health and public safety. With the increase in alcohol production and consumption annually, public health problems caused by alcohol abuse are becoming increasingly prominent.

Alcohol can induce damage to the central nervous system, leading to changes in the brain function and structure⁶. Studies have reported that cognitive impairment in patients with alcohol dependence is closely related to white matter atrophy, which affects vision, space, memory, attention, and some other functions, such as blood flow⁷. Pandey et al. reported that executive dysfunction in patients with alcohol dependence was associated with prefrontal lobe damage, whereas memory dysfunction was associated with the hippocampal volume⁸. Findings from previous reports have suggested that most patients with alcohol dependence have different degrees of cerebral hypoperfusion, including that in the cerebral cortex and subcutaneous structural regions, but most researchers believe that the prefrontal cortex is the core region⁹. Oxygenated hemoglobin (oxy-Hb) is a substance produced by the reversible combination of hemoglobin and oxygen molecules, which helps transport oxygen¹⁰. The blood-oxygen level has been shown to rapidly rise in the human brain engaged in cognitive activities through the action of the neurovascular matching mechanism, which significantly increases the concentration of oxy-Hb in the cognitive activity area of the brain¹¹. Moreover, a study that included patients with schizophrenia showed that the concentration of oxy-Hb in the prefrontal lobe was significantly reduced during cognitive activity when compared to healthy individuals¹².

In the past, findings from several studies have revealed the influence of oxy-Hb levels on the cognitive impairment in patients with alcohol dependence. The aim of this present study was to evaluate the diagnostic value of oxy-Hb in the cognitive impairment of patients with alcohol dependence by measuring the level of oxy-Hb in the prefrontal lobe of patients, thus providing insights into clinical interventions for such patients.

Materials and methods

STUDY POPULATION

Seventy-one male patients with alcohol dependence hospitalized in Hengshui Seventh People's Hospital were selected for the observation group. The inclusion criteria for patients with alcohol dependence were as follows: 1) the diagnosis of the patients should conform to the diagnostic criteria of alcohol dependence in the International Classification of Diseases-10; 2) males age from 30 to 60 years; 3) patients with normal intelligence; 4) patients who stopped drinking alcohol at least 14 days prior to the test and also stopped using alternative sedatives. The exclusion criteria were as follows: 1) patients with organic mental disorders; 2) patients with a history of central nervous system disease, such as stroke, epilepsy, and tumor; 3) patients with a history of serious chronic diseases, such as hypertension and diabetes. Additionally, 70 healthy adult male patients were selected during the same period for the control group. The study was approved by the Ethics Committee of Hengshui Seventh People's Hospital, and all the members who participated in the study had signed the informed consent form after being fully informed.

COLLECTION OF SAMPLE AND MEASUREMENT OF OXY-Hb

In this study, the oxy-Hb content was determined using the ETG-4000 type quantitative imaging device of brain function developed and manufactured by Hitachi. This quantitative imaging device system has 17 optical signal transmitters that emit infrared light of 695 nm and 830 nm and measure the cognitive activities in the subjects' brains during the stimulus tasks to measure the oxy-Hb levels. The device captures infrared signals through a detector in which the sensor is attached to a soft and flexible cloth cover, and the cloth cover is fixed on the forehead of the subject to examine the frontal lobe of the brain. The oxy-Hb level was measured using this imaging device, and the data were analyzed using the VerL.61e system analysis software attached to the test instrument.

COGNITIVE FUNCTION ASSESSMENT

Two technicians specializing in psychological assessment evaluated the patients with alcohol dependence who stopped taking antipsychotic medications. Subjects were required to complete the Wisconsin card sorting test (WCST), brief visuospatial memory test-revised (BVM-T-R), digital sequence test, digital span test, and continuous performance test (CPT).

The subjects completed the WCST after selecting six correct categories. During the test, subjects were unaware of the category principle, and the system automatically informed the subjects of the results. The responses administered, correct responses, categories completed, trials to compete first category, perseverative responses, and perseverative response errors of subjects were recorded. BVM-T-R primarily measured the memory ability of subjects. The total score was the sum of the scores of three tests. The higher the total score, the stronger the memory ability of the subjects. Similarly, the digital sequence test measured the operating memory ability of subjects. The test took approximately 5 min to complete, with 1 and 0 points awarded for correct and wrong responses, respectively. A higher total score corresponded to a higher operational memory ability. Meanwhile, the digital span test measured the attention and short-term memory of subjects. A higher score indicated a higher level of attention. Further, CPT was used to examine the sustained attention level of subjects, which is the "maintenance factor" of attention and is related to the level of arousal. Reaction time, reaction time change index, number of mistakes and omissions of subjects were recorded, and a subject with a higher score was considered to have a higher level of attention.

DATA ANALYSIS

SPSS 22.0 was used for data processing. The quantitative data followed the normal distribution and were represented by mean \pm standard deviation ($\bar{x} \pm SD$). The independent sample *t*-test was used for comparison between two groups. Moreover, the results of the correlation study were analyzed using the Pearson correlation coefficient method. Further, the receiver operating characteristic (ROC) curve was used to evaluate the diagnostic value of oxy-Hb in alcohol-dependent cognitive dysfunction. $P < 0.05$ indicated that the difference was statistically significant.

Results

OXY-HB CONTENT IN PATIENTS WITH ALCOHOL DEPENDENCE

A total of 141 subjects were tested, which included 70 healthy subjects and 71 patients with alcohol de-

pendence. No statistically significant differences were observed between the two groups in terms of age, BMI, smoking history, and education level; hence, the groups were comparable (table 1), ($p > 0.05$). The results revealed a significant declining trend in the levels of oxy-Hb in the brains of patients with alcohol dependence as compared to that in healthy individuals (Figure 1), ($p < 0.001$).

COMPARISON OF COGNITIVE FUNCTION SCORES BETWEEN THE OBSERVATION AND CONTROL GROUPS

As shown in table 2, the scores of responses administered, trials to compete first category, perseverative responses, and perseverative response errors

Table 1. Clinical data of the study population.

Variables	All subjects (n=141)		P value
	Healthy individuals (n=70)	Alcohol dependence (n=71)	
Age	47.33 \pm 7.77	49.14 \pm 8.14	0.177
BMI	23.51 \pm 2.84	23.54 \pm 2.97	0.959
Smoking status (n)			0.052
Yes	28	40	
No	42	31	
Education Level (n)			0.271
Junior high and below	30	37	
Junior high school above	40	34	

Legend: BMI= body mass index.

Data are expressed as n or mean \pm standard deviation.




Figure 1. The concentration of oxygenated hemoglobin (oxy-Hb) in the frontal lobe of patients with alcohol dependence decreased compared to that in healthy controls.

*** $P < 0.001$.

Table 2. Comparison of cognitive function test scores between groups.

Variables	All subjects (N=141)		P value
	Healthy individuals (n=70)	Alcohol dependence (n=71)	
Executive function			
Response administered	106.73±8.91	139.91±17.43	<0.001
Correct responses	71.47±6.19	46.31±6.54	<0.001
Categories completed	4.81±0.80	1.58±0.62	<0.001
Trials to compete first category	24.42±5.68	58.28±10.65	<0.001
Perseverative responses	26.24±5.09	38.22±8.35	<0.001
Perseverative responses errors	20.79±6.84	34.79±6.18	<0.001
Memory function			
Brief visuospatial memory test-revised	29.00±1.69	18.30±2.97	<0.001
Digital sequence test	27.40±2.22	18.31±3.00	<0.001
Attention function			
Digital span test	20.89±2.22	14.89±1.78	<0.001
Continuous performance test	3.00±0.80	1.75±0.68	<0.001

were higher, whereas the scores of correct responses and categories completed were lower in the observation group as compared to that in control group ($p<0.001$). Moreover, the scores of memory function (BVMT-R, digital sequence test) and attention function (digital span test, CPT) were significantly lower in the observation group than in the control group ($p<0.001$).

CORRELATION BETWEEN THE OXY-Hb LEVEL AND COGNITIVE FUNCTION SCORES IN PATIENTS WITH ALCOHOL DEPENDENCE

Pearson correlation analysis has shown that the scores of correct responses, categories completed, BVMT-R, digital sequence test, digital span test, and CPT were positively correlated with the levels of oxy-Hb (table 3), ($p<0.05$). However, the scores of administered, trials to compete first category, and perseverative response errors were negatively correlated with

Table 3. Correlation between relative concentration of Oxy-Hb and various indicators.

Variables	Alcohol dependence group (n=71)	
	r	p
Executive function		
Response administered	-0.749	<0.001
Correct responses	0.401	0.001
Categories completed	0.293	0.013
Trials to compete first category	-0.827	<0.001
Perseverative responses	-0.193	0.108
Perseverative response errors	-0.693	<0.001
Memory function		
Brief visuospatial memory test-revised	0.601	<0.001
Digital sequence test	0.333	0.005
Attention function		
Digital span test	0.577	<0.001
Continuous performance test	0.494	<0.001

the oxy-Hb levels ($p<0.001$). Besides, no correlation was observed between the score of perseverative response and oxy-Hb level ($p=0.108$).

DIAGNOSTIC VALUE OF OXY-Hb IN COGNITIVE IMPAIRMENT IN PATIENTS WITH ALCOHOL DEPENDENCE

Patients in the observation group received MMSE and MoCA tests and were categorized into cognitive impairment and non-cognitive impairment groups according to the results (results shown in table 4). As illustrated in figure 2A, the oxy-Hb levels in patients with cognitive impairment were lower than those in the non-cognitive impairment group ($p<0.001$). Subsequently, the oxy-Hb in the observation group was analyzed by ROC analysis. As shown in figure 2B, the area under ROC curve of the oxy-Hb level in the observation group was 0.930 at the cut-off value of 0.135, and 95% CI was 0.874-0.987. The sensitivity and specificity of oxy-Hb in the diagnosis of alcohol-dependent cognitive dysfunction were 89.1% and 84.0% respectively.

Discussion and conclusions

Cognition is the process in which the individual's sensory organs process external information input and convert it into an individual's internal psychological activities¹³. These activities mainly include

Table 4. Comparison of MMSE and MoCA scores between two groups.

Variables	Alcohol-dependent patients (n = 71)		P value
	Poor cognitive function (n = 22)	Good cognitive function (n = 49)	
MMSE score	21.36 ± 1.62	25.75 ± 1.60	< 0.001
MoCA score	20.45 ± 1.92	27.55 ± 1.19	< 0.001

Legend: MMSE= Mini-Mental State Examination; MoCA= Montreal Cognitive Assessment. Data are expressed as n or mean ± standard deviation.

perception, memory, attention, thinking, language, and orientation. The factors that may directly or indirectly cause structural and functional damage to the cerebral cortex can cause cognitive dysfunction in individuals through different mechanisms that include chronic brain tissue injury, chronic systemic diseases, mental and psychological abnormalities, and other humanistic factors^{14,15}. Cognitive dysfunction adversely affects different aspects of an individual's regular life, including study, work, and social relationships. The mechanism underlying cognitive impairment in patients with alcohol dependence has

not yet been clarified, but it is considered to be related to the central nervous system damage caused by the neurotoxic effect of alcohol. In humans, the prefrontal cortex is primarily involved in advanced cognitive functions, such as perception, memory, execution, decision-making, attention, and reasoning¹⁶. By measuring the change in the oxy-Hb concentration in the prefrontal cortex during cognitive activities, we can preliminarily determine whether the cerebral hemodynamics of subjects is normal or whether there is the possibility of damage.

Alcohol is an inhibitor of the central nervous system and exerts a direct toxic effect on the nervous system, which can alter brain function and structure^{17,18}. Studies have shown that patients with alcohol dependence show varying degrees of decreased cerebral blood perfusion and cerebral blood flow in the prefrontal lobe¹⁹. These pathological factors may cause diffuse cerebral blood supply insufficiency, severe brain hypoxia, or even brain atrophy, which may directly induce cognitive dysfunction in patients with alcohol dependence²⁰. During the cognitive activity of the brain, oxygen consumption and blood perfusion in the brain tissue increase, which upregulates the concentration of oxy-Hb in the active area of the brain tissue²¹. In this study, the oxy-Hb concentration in the prefrontal lobe of patients with alcohol dependence and healthy controls during cognitive activities was measured using a brain function imaging in-




Figure 2. (A) The concentration of oxy-Hb in patients with poor cognitive function was lower than that in individuals with good cognitive function group; (B) Receiver operating characteristic (ROC) curve analysis. The area under the curve of oxy-Hb was 0.926, and the sensitivity and specificity were 88.7% and 82.9%, respectively. **P < 0.001.

strument. The results showed that the concentration of oxy-Hb in patients with alcohol dependence was significantly lower than that in healthy individuals. Besides, these results were consistent with findings reported by Obata et al., which demonstrated that the oxy-Hb concentration was lower in individuals after alcohol consumption²². Therefore, we concluded that the phenomenon is related to the neuroinhibitory effect of alcohol. Patients with alcohol dependence are conditioned to long-term alcohol consumption, which exerts a neurotoxic effect on the brain, leading to functional and structural changes in the central nervous system and causing cerebral blood flow hypoperfusion, cerebral blood oxygen saturation reduction, and long-term hypoxia in the brain²³. In patients with alcohol dependence, sufficient oxygen is unavailable for neuronal activities during cognitive activities, and thus, oxygen consumption in the relevant brain regions activated by cognitive activities decreases, thereby reducing the relative concentration of oxy-Hb in the brain regions.

It is well established that alcohol affects human cognitive function. Rao et al. found that memory function in patients with alcohol dependence was severely impaired, with the significant impairment of transient and short-term memory²⁴. Findings from several studies have confirmed that cognitive dysfunction related to mental illness is closely correlated with the relative concentration of oxy-Hb in the prefrontal lobe²⁵. Pu et al. showed that the prefrontal blood flow and the relative concentration of oxy-Hb decreased significantly in patients with depression during the onset of the disease²⁶. In this study, the executive function, memory function, and attention function in the observation group declined compared to that in the control, suggesting that patients with alcohol dependence showed impaired cognitive function. The results of Pearson's correlation analysis showed that the scores of correct responses, categories completed, BVMT-R, digital sequence test, digital span test, and CPT were positively correlated with the oxy-Hb concentration. Besides, the scores of responses administered, trials to compete first category, and perseverative response errors were negatively correlated with the oxy-Hb concentration. In addition, the concentration of oxy-Hb in the observation group was analyzed by the ROC curve, and the results showed that the oxy-Hb had high accuracy in the diagnosis of cognitive impairment caused by alcohol dependence.

The present study had certain limitations, including the fact that this was a single-center study with small sample size. Moreover, the years of alcohol dependence of patients were not included in the study, which may have led to a certain degree of difference in the brain function and structure as well as changes in the psychological behavior of patients. Besides, the

evaluation of cognitive functions in patients with alcohol dependence was not comprehensive, and several other cognitive functions, such as decision-making, language, comprehension, judgment, and calculation, were not evaluated. Thus, a follow-up study with a large sample size is warranted.

In conclusion, we demonstrated that cognitive impairment occurs in patients with alcohol dependence, and the concentration of oxy-Hb in the prefrontal cortex of patients decreases compared to that in healthy people. This indicates that oxy-Hb detection can be a preliminarily determinant to suggest whether the cognitive function of patients with alcohol dependence is impaired.

Conflict of interests: the authors have no conflict of interests to declare.

References

1. Roos CR, Maisto SA, Witkiewitz K. Coping mediates the effects of cognitive-behavioral therapy for alcohol use disorder among out-patient clients in Project MATCH when dependence severity is high. *Addiction* 2017; 112: 1547-57.
2. Affan RO, Huang S, Cruz SM, Holcomb LA, Nguyen E, Marinkovic K. High-intensity binge drinking is associated with alterations in spontaneous neural oscillations in young adults. *Alcohol* 2018; 70: 51-60.
3. Kim KM, Kim JS, Kim GJ, et al. The readiness to change and insight in alcohol dependent patients. *J Korean Med Sci* 2007; 22: 453-8.
4. Park JH, Kim SG, Kim JH, Lee JS, Jung WY, Kim HK. Spicy food preference and risk for alcohol dependence in Korean. *Psychiatry Investig* 2017; 14: 825-9.
5. Satou R, Oka S, Sugihara N. Risk assessment of fluoride daily intake from preference beverage. *J Dent Sci* 2021; 16: 220-8.
6. Zhang X, Sun L, Wang L, et al. The effects of histone deacetylase inhibitors on the attentional set-shifting task performance of alcohol-dependent rats. *Brain Res Bull* 2019; 149: 208-15.
7. Clarke TK, Smith AH, Gelernter J, et al. Polygenic risk for alcohol dependence associates with alcohol consumption, cognitive function and social deprivation in a population-based cohort. *Addict Biol* 2016; 21: 469-80.
8. Pandey AK, Ardekani BA, Kamarajan C, et al. Lower prefrontal and hippocampal volume and diffusion tensor imaging differences reflect structural and functional abnormalities in abstinent individuals with alcohol use disorder. *Alcohol Clin Exp Res* 2018; 42: 1883-96.
9. Zuo N, Fang J, Lv X, et al. White matter abnormalities in major depression: a tract-based spatial statistics and rumination study. *PLoS One* 2012; 7: e37561.
10. Takemura N, Sakatani K, Yoshino A, Hirayama T, Katayama Y. Physiological mechanism of increase in deoxyhemoglobin concentration during neuronal activation in patients with cerebral ischemia: a simulation study with the balloon model. *Adv Exp Med Biol* 2014; 812: 225-31.
11. De la Torre GG. Cognitive neuroscience in space. *Life (Basel)* 2014; 4: 281-94.
12. Chou PH, Koike S, Nishimura Y, et al. Similar age-related decline in cortical activity over frontotemporal regions in schizophrenia: a multichannel near-infrared spectroscopy study. *Schizophr Bull* 2015; 41: 268-79.
13. DeLong KA, Troyer M, Kutas M. Pre-processing in sen-

- tence comprehension: Sensitivity to likely upcoming meaning and structure. *Lang Linguist Compass* 2014; 8: 631-45.
14. Nestor LJ, Paterson LM, Murphy A, et al.; ICCAM Consortium. Naltrexone differentially modulates the neural correlates of motor impulse control in abstinent alcohol-dependent and polysubstance-dependent individuals. *Eur J Neurosci* 2019; 50: 2311-21.
 15. Andreassen Ø, Brønnick K, Njå AL, Furulund E, Nesvåg S. The effect of high-intensity interval/circuit training on cognitive functioning and quality of life during recovery from substance abuse disorder. A study protocol. *Front Psychol* 2019; 10: 2564.
 16. Crowe SF, Cammisuli DM, Stranks EK. Widespread cognitive deficits in alcoholism persistent following prolonged abstinence: an updated meta-analysis of studies that used standardised neuropsychological assessment tools. *Arch Clin Neuropsychol* 2019; 35: 31-45.
 17. Rao X, Thapa KS, Chen AB, et al. Allele-specific expression and high-throughput reporter assay reveal functional genetic variants associated with alcohol use disorders. *Mol Psychiatry* 2021; 26: 1142-51.
 18. Chen MM, Palmer JL, Ippolito JA, Curtis BJ, Choudhry MA, Kovacs EJ. Intoxication by intraperitoneal injection or oral gavage equally potentiates postburn organ damage and inflammation. *Mediators Inflamm* 2013; 2013: 971481.
 19. Fu C, Shi D, Gao Y, Xu J. Functional assessment of prefrontal lobes in patients with major depression disorder using a dual-mode technique of 3D-arterial spin labeling and (18)F-fluorodeoxyglucose positron emission tomography/computed tomography. *Exp Ther Med* 2017; 14: 1058-64.
 20. Bhowmik A, Haldar SK, Alon OE. Impact of the transverse direction on the many-body tunneling dynamics in a two-dimensional bosonic Josephson junction. *Sci Rep* 2020; 10: 21476.
 21. Diaz-Parra A, Perez-Ramirez U, Pacheco-Torres J, et al. Evaluating network brain connectivity in alcohol post-dependent state using Network-Based Statistic. *Annu Int Conf IEEE Eng Med Biol Soc* 2017; 2017: 533-6.
 22. Obata A, Morimoto K, Sato H, et al. Acute effects of alcohol on hemodynamic changes during visual stimulation assessed using 24-channel near-infrared spectroscopy. *Psychiatry Res* 2003; 123: 145-52.
 23. Le Berre AP, Müller-Oehring EM, Schulte T, Serventi MR, Pfefferbaum A, Sullivan EV. Deviant functional activation and connectivity of the right insula are associated with lack of awareness of episodic memory impairment in nonamnesic alcoholism. *Cortex* 2017; 95: 15-28.
 24. Wang GJ, Shokri Kojori E, Yuan K, et al. Inhibition of food craving is a metabolically active process in the brain in obese men. *Int J Obes (Lond)* 2020; 44: 590-600.
 25. Ehlis AC, Schneider S, Dresler T, Fallgatter AJ. Application of functional near-infrared spectroscopy in psychiatry. *Neuroimage* 2014; 85 Pt 1: 478-88.
 26. Pu S, Nakagome K, Yamada T, et al. Prefrontal activation predicts social functioning improvement after initial treatment in late-onset depression. *J Psychiatr Res* 2015; 62: 62-70.

Corresponding author:
Dr. Xiangyang Zheng
Department of Psychiatry
Hengshui Seventh People's Hospital
No.1769 Shengli West Road
Hengshui 053000, China
E-mail: zhengxiangyang05@163.com