

Perinatal mental health and Covid-19-related distress: the role of personality traits

MELANIA MARTUCCI^{1*}, MATTEO PANFILI^{1*}, NICOLETTA GIACCHETTI¹, FRANCESCO SAVERIO BERSANI¹, PAOLA CIOLLI², ALESSANDRA FORTE¹, CARLA SOGOS¹, FRANCA ACETI¹

¹Department of Human Neurosciences, Sapienza University of Rome, Italy; ²Department of Gynecology, Obstetrics and Urology, Sapienza University, Rome, Italy.

*Contributed equally.

Summary. Background. Perinatal mental health is a topic of growing interest, that could affect mothers in a period of high vulnerability, and the impact of coronavirus disease 2019 (Covid-19) pandemic is an important factor to consider in this field. The aim of our study is to study the correlations between five dimensions of personality and subjective Covid-19-related distress in a sample of women in the perinatal period. **Methods.** The study included 114 Italian women in the perinatal period. Subjects were asked to complete the Big Five Inventory (BIG-5) and a version of the Impact of Event Scale – Revised (IES-R) anchored to Covid-19-related distress. **Results.** When the BIG-5 personality traits and several confounding variables were included in a regression model with IES-R total score as criterion, neuroticism subscale of BIG-5 inventory was the only variable independently associated with higher IES-R total score ($p < 0.001$). **Conclusion.** Our study highlights the importance of considering the personality vulnerability factors that can worsen psychopathological symptoms of women in the perinatal period, especially in periods of high psychological stress.

Key words. Covid-19, perinatal mental health, personality.

La salute mentale nel periodo perinatale e lo stress correlato al Covid-19: il ruolo dei tratti di personalità.

Riassunto. Introduzione. La salute mentale perinatale è un argomento di crescente interesse, che riguarda le madri in un periodo di elevata vulnerabilità. L'impatto della pandemia da Covid-19 è un fattore importante da considerare in questo campo. Lo scopo del nostro studio è studiare le correlazioni tra le cinque dimensioni della personalità e lo stress percepito correlato al Covid-19 in un campione di donne nel periodo perinatale. **Metodi.** Lo studio ha incluso 114 donne italiane nel periodo perinatale. Ai soggetti è stato chiesto di completare il Big Five Inventory (BIG-5) e una versione della Impact of Event Scale – Revised (IES-R) per valutare lo stress correlato al Covid-19. **Risultati.** È stato effettuato un modello di regressione lineare includendo i tratti di personalità valutati mediante BIG-5 e diverse variabili confondenti; la sottoscala nevroticismo del questionario BIG-5 è risultata l'unica variabile indipendente associata a un punteggio totale IES-R maggiormente elevato ($p < 0,001$). **Conclusione.** Il nostro studio evidenzia l'importanza di considerare i fattori di vulnerabilità della personalità che possono peggiorare i sintomi psicopatologici delle donne nel periodo perinatale, soprattutto nei periodi di elevato stress psicologico.

Parole chiave. Covid-19, personalità, salute mentale perinatale.

Introduction

Perinatal mental health is a topic of growing interest, with distinct clinical conditions that could affect mothers in a period of high vulnerability. The peripartum is a period of special needs for women with pre-existing psychopathological conditions such as Major Depressive Disorders (MDD), Borderline Personality Disorders (BPD), Post-Traumatic Stress Disorder (PTSD) and other psychiatric disorders, who are at higher risk of relapses^{1,2}. It is also a time of neuroendocrine, biological, psychological and social adaptation, hence new mental disorders may appear, such as perinatal depression and puerperal psychosis^{1,2}.

Evidence estimates that the prevalence rate of perinatal mood and anxiety disorders range approxi-

mately between 10 and 20%^{3,4}. Antenatal depression and anxiety, stressful life events, poor relationship quality and social support, lower socioeconomic status, single marital status and unintended pregnancy are factors which have showed associations with increased risk of depressive symptoms during pregnancy⁵. A recent review suggested that the strongest risk factors for postpartum depression (PD) are antenatal depression and domestic abuse⁶.

In some cases, depressive symptoms hide a personality disorder. In fact, there are numerous evidence that report perinatal depressive symptoms in patients with obsessive-compulsive, dependent, avoidant and borderline personality disorders. Moreover, some personality traits have been considered as risk factors for perinatal depression: dependency⁷, obsession⁸, neuroticism⁹, the tendency to severe self-criticism¹⁰.

The impact of the pandemic of coronavirus disease 2019 (Covid-19) has been investigated as another important factor to consider in this field. Previous studies have shown that patients in the prenatal and postnatal periods are particularly at risk of developing mental disorders during health or social disasters¹¹. Thus, the prevalence of perinatal anxiety and depression has significantly raised during the pandemic. The results of a systematic review including 11,187 pregnant women in China showed that the prevalence of anxiety was 34% and the prevalence of both anxiety and depression was 18%¹². The factors which may have contributed to the increase of perinatal mental disorders are the fear of being infected or infected others (intrauterine transmission, passage of the virus from mother to baby during childbirth, infection through breast-feeding), and the uncertainty about the effects of the virus on the fetuses and infants. It is important to highlight that public health policies such as lockdown, limiting prenatal visits, social distancing measures, and their many associated socio-economic consequences (unemployment, loss of income, and domestic violence) may have been an additional challenge for perinatal mental health¹³.

Moreover, during the pandemic, anti-vaccine positions have increased, lingering doubts about vaccine safety. These doubts were particularly important for pregnant women fearing effects on their fetus¹³. Prenatal exposure to negative maternal psychological factors, including stress due to disasters, can be associated with a range of poor outcomes for the offspring, including preterm birth, adverse obstetrical outcomes (such as preeclampsia), and low birth-weight¹⁴. In addition, pregnant women with Covid-19 are at risk for placental dysfunction, characterized by inflammatory changes, massive perivillous fibrin depositions and placental necrosis, and related fetal distress, specifically when infected with the Delta variant¹⁵. It is also known that certain maternal infections during pregnancy may have a role in the development of a range of mental disorders in the offspring, such as schizophrenia, autism spectrum disorders, and attention deficit/hyperactivity disorder¹⁶⁻¹⁸.

Even though the Covid-19 public health emergency has been declared ended, it is important to increase knowledge about perinatal mental disorders by studying the implications of such period of high distress on perinatal mental health. Thus, the aim of our study has been to study the relationships of specific dimensions of personality with the degree of subjective distress caused by Covid-19 pandemic in a sample of Italian women in the perinatal period.

Materials and methods

PARTICIPANTS

The sample was made of 114 Italian women of 32.06 ± 4.96 years (age range: 20-46); 57 of them (50%) were pregnant, 57 (50%) had given birth within the previous 12 months.

Such amount of participants was selected as a sample size calculation performed with the G*Power 3.1 software¹⁹ indicated that 92 subjects are needed to achieve a moderate effect size ($f^2=0.15$) with power=0.80 and alpha=0.05 in a regression analysis with five tested predictor and 12 total number of predictors (details on predictors are given below).

The recruitment was performed between April 2020 and January 2021. Among the participants, 35 (24.3%) were recruited from obstetrics department, 9 (7.8%) from the outpatients gynecologic service, 70 (61.4%) from the general population (i.e. they were volunteers).

Inclusion criteria were being pregnant or having a child under the age of 1. The exclusion criteria were as follows: the refusal to provide an informed consent, less than 18 years of age, the presence of a diagnosis of mental retardation or schizophrenia spectrum disorder, a poor knowledge of the Italian language or other limitation in verbal communication which compromised the patient's ability to perform the research. Subjects voluntarily participated in the study, i.e. they did not receive payment or other compensation.

MATERIALS

All participants were administered a checklist assessing certain clinical and socio-demographic information: age, pregnancy status, education level, ongoing psychotropic medications, previous pregnancies, intended pregnancy, family history of mental disorders.

Subjects were asked to complete the self-report measures Big Five Inventory (BIG-5)²⁰ and Impact of Event Scale - Revised (IES-R)²¹.

The BIG-5 Inventory is a 44-item inventory that measures an individual on the Big Five Factors (dimensions) of personality²⁰. The items of this scale were selected from Big Five prototype definitions that had been developed through expert ratings to allow efficient and flexible assessment of the five personality dimensions (extraversion, agreeableness, conscientiousness, neuroticism, and openness)²⁰. In the present sample, Cronbach's alpha for each subscale was: 0.726 for extraversion, 0.673 for agreeableness, 0.716 for conscientiousness, 0.800 for neuroticism and 0.745 for openness.

The IES-R scale, is a self-report questionnaire of 22 items, appropriate to measure the subjective

response to a specific traumatic event in the adult population, especially in the response sets of intrusion (intrusive thoughts, nightmares, intrusive feelings and imagery, dissociative-like re-experiencing), avoidance (numbing of responsiveness, avoidance of feelings, situations, and ideas), and hyperarousal (anger, irritability, hypervigilance, difficulty concentrating, heightened startle)²¹. In the present study, IES-R was specifically anchored to Covid-19-related distress; IES-R total score was used, and Cronbach's alpha was 0.917.

STATISTICAL ANALYSES

All analyses were performed using the SPSS (26.0) statistical package (IBM, Armonk, NY, USA). Descriptive analyses with means and standard deviations were performed. Spearman correlation coefficients (ρ) were calculated to study the associations of the explored variables. A linear regression was performed with those BIG-5 subscales which were significantly associated with IES by Spearman correlation as the determinants, and IES total score as outcome (dependent variable). The following variables were also included in the regression model as covariates that could be considered as potential confounders of the association between personality traits and psychopathological reaction to stress: age, pregnancy status, education, use of psychiatric medication at the time of evaluation, previous pregnancies, planned pregnancy or not, family history of psychiatric disorders. The major assumptions of multiple regression (zero conditional mean of errors, homoscedasticity, normal distribution of errors, multicollinearity, outliers) were checked.

Results

The mean age of the present sample of women was 32.06 ± 3.70 years. Among pregnant women week of pregnancy was 29.56 ± 8.33 , among women in their post-partum period the age of the offspring was 4.38 ± 2.3 weeks. 70.2% of the pregnancies were intended, 29.8 % were not. 60.5% of the women were at their first pregnancy, 39.5% were multiparous; 7.4% of the women had lower secondary education diploma, 53.7% had upper secondary education diploma, and 39.5 % had a Bachelor and Master Degree; 25.4% of the women had a family history of psychiatric disorders, and 6.1% of them was taking psychotropic medications. Further details on descriptive statistics are reported in Table 1.

Spearman correlations showed that IES-R total score was significantly associated with the BIG-5 subscales on neuroticism ($\rho=0.379$; $p<0.001$), extraversion ($\rho=-0.221$; $p=0.018$), conscientious-

Table 1. Descriptive statistics of the sample (n=114, 57 pregnant women, 57 women who had given birth in the previous 12 months).

Variables	
Age, M \pm SD	32.06 \pm 3.70
Week of pregnancy (among pregnant women, n=57), M \pm SD	29.56 \pm 8.33
Weeks of age of the newborn (among women in the postnatal period), M \pm SD	4.38 \pm 2.3
Education	
Lower Secondary Education Diploma, N (%)	7 (7.4)
Upper Secondary Education Diploma, N (%)	63 (53.7)
Bachelor and Master Degree, N (%)	44 (39.5)
Intended pregnancy, N (%)	80 (70.2%)
Primiparous, N (%)	69 (60.5%)
Presence of family history of psychiatric disorders, N (%)	29 (25.4%)
Ongoing psychotropic medications, N (%)	7 (6.1%)

ness ($\rho=-0.196$; $p=0.036$), and openness ($\rho=-0.195$; $p=0.038$), while it was not significantly associated with the agreeableness subscale.

The linear regression model with IES total score as criterion, with extraversion, conscientiousness, neuroticism, and openness as determinants, and with age, pregnancy status, education, use of psychiatric medication at the time of evaluation, previous pregnancies, planned pregnancy or not, family history of psychiatric disorders as covariates, explained 19.9% of the variability of the data ($R^2=0.199$, adjusted $R^2=0.112$). In this model, neuroticism was significantly ($\beta=0.375$, $p<0.001$) independently associated with IES, so that a more pronounced form of such personality pattern was associated with higher IES score even when controlling for the presence of the other variables. Further details are reported in table 2 and in figure 1.

Discussion

Our study showed that neuroticism was the unique personality dimension detected by BIG-5 Inventory significantly and independently correlated with the perceived impact of the pandemic, also when certain potential confounding factors were controlled for.

Literature reported that neuroticism refers to individual differences in negative emotionality and predicts a range of negative outcomes, including low subjective well-being²² and poor physical and mental health²³. Prior research has established that personality differences modulate the stress process²⁴.

Table 2. Linear regression with Impact of Event Scale – Revised total score as criterion.

	Standardized Coefficient (B)	P	95% CI	
			Lower Bound	Upper Bound
Age	-0.058	0.579	-0.041	0.023
Pregnancy	0.065	0.489	-0.183	0.380
Education	0.008	0.940	-0.213	0.229
Ongoing psychotropic medications	-0.071	0.507	-0.894	0.445
Previous pregnancies	-0.036	0.705	-0.348	0.236
Intended pregnancy	0.019	0.838	-0.274	0.338
Family history of psychiatric disorders	-0.164	0.094	-0.625	0.050
Conscientiousness	-0.083	0.396	-0.363	0.145
Openness	0.026	0.800	-0.219	0.283
Extraversion	0.015	0.898	-0.268	0.305
Neuroticism	0.375	<0.001*	0.167	0.624

*p<0.05




Figure 1. Regression scatterplot showing the significant association between IES-R total score and neuroticism BIG-5 subscale, after controlling for confounding factors (age, pregnancy, education, ongoing psychotropic medications, previous pregnancies, intended pregnancy, family history of psychiatric disorders, conscientiousness, openness and extraversion BIG-5 subscales).

More specifically, neuroticism, is characterized by depression, anxiety and greater reactivity to stressful events²⁴. Extraversion, instead, characterized by sociability, assertiveness, and energy, has been typically associated with lower stress reactivity and lower negative affect²⁴. However, the transactional theory of stress suggests that personal characteristics will not necessarily predict stress reactivity in all circumstances, but may function differently when the situational context varies²⁵.

Regarding Covid-19 pandemic, evidence suggests a potential function of neuroticism in aggravating the stress process. Bellingtier et al.²⁵ reported that in-

dividuals higher in neuroticism were more likely to perceive the pandemic as strongly affecting their personal lives with greater perceived stress and worry, higher levels of depression and anxiety symptoms and lower well-being. Hence, the feelings of insecurity, instability, difficulty in controlling emotions, difficulty in managing interpersonal dynamics peculiar to this dimension may play a relevant role in the psychic discomfort resulting from an event of this magnitude.

A recent study reports some relevant data regarding neuroticism and perceived stress. It suggests that highly neurotic individuals were at higher risk for

stress during the Covid-19 pandemic. In addition, neuroticism significantly mediated the relationship between fear of Covid-19 and perceived stress. More specifically, higher levels of neuroticism were related to higher levels of fear of Covid-19 and perceived stress²⁶. Hence, emotional vulnerability related to pregnancy or the recent birth of a child, can contribute to making women with this personality trait particularly at risk in terms of perinatal mental health²⁴.

Exploring mental health status of women in the perinatal period is an important issue, also in light of the recent evidence about maternal mental health and child development. In fact, numerous studies showed that the children of mothers with perinatal mental disorders had a higher risk, compared to controls, for the early onset of emotional development disorders and lower emotion recognition at ten years^{27,28}, possible neuropsychological or cognitive deficits²⁹⁻³¹, internalizing and externalizing behavioral disorders²⁸, sleep disturbances^{32,33}, persistent lower growth³⁴ and long-term consequences on the child and adolescent mental health³⁵.

Limits and future perspectives

The results of the present research should be interpreted taking into consideration certain limitations. This is a cross-sectional study, therefore the existence of a causal link between the variables that are associated cannot be stated with certainty, nor the possible implication of this potential causal link. A longitudinal study of the variables involved, aimed to evaluate their evolution pre and post pandemic could provide useful information. Although the neuroticism subscale of the BIG5 and the IES-S scale evaluate different psychological constructs, there might be some degree of similarity in certain items of the two scales, which could contribute to explain the observed association. Moreover, a critical point is the small clinical sample, it should be expanded to make the results generalizable.

Overall, our study contributes to highlight potential implications of Covid-19 pandemic on perinatal mental health, also considering the personality vulnerability factors. It is important to detect these psychological situations in order to take care of the mothers, to prevent the consequences on childbirth and to increase knowledge in this field of research³⁵⁻⁴³.

Funding: this research received no external funding.

Institutional review board statement: the study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of Sapienza University of Rome (approval date 31 January 2018, approval date 4816).

Informed consent statement: informed consent was obtained from all subjects involved in the study.

Conflicts of interests: the authors have no conflict of interests to declare.

References

- O'Hara MW, Swain A. Rates and risk of post-partum depression: a meta-analysis. *Int Rev Psychiatry* 1996; 8: 37Y54.
- Viguera AC, Cohen LS, Baldessarini RJ, Nonacs R. Managing bipolar disorder during pregnancy: Weighing the risks and benefits. *Can J Psychiatry* 2002; 47: 426Y436.
- Fawcett EJ, Fairbrother N, Cox ML, White IR, Fawcett JM. The prevalence of anxiety disorders during pregnancy and the postpartum period: a multivariate bayesian meta-analysis. *J Clin Psychiatry* 2019; 80: 18r12527.
- Howard LM, Khalifeh H. Perinatal mental health: a review of progress and challenges. *World Psychiatry* 2020; 19: 313-27.
- Lancaster CA, Gold KJ, Flynn HA, Yoo H, Marcus SM, Davis MM. Risk factors for depressive symptoms during pregnancy: a systematic review. *Am J Obs Gynecol* 2010; 202: 5-14.
- Hutchens BF, Kearney J. Risk factors for postpartum depression: an umbrella review. *J Midwifery Womens Health* 2020; 65: 96-108.
- Priel B, Besser A. Vulnerability to postpartum depressive symptomatology: dependency, self-criticism, and the moderating role of antenatal attachment. *J Soc Clin Psychol Meas* 1999; 18: 240-53.
- Akman I, Kuscu MK, Yurdakul Z, et al. Breastfeeding duration and postpartum psychological adjustment: role of maternal attachment styles. *J Paediatr Child Health* 2008; 44: 369-73.
- Podolska MZ, Bidzan M, Majkovicz M, Podolski J, Sipak-Szmigiel O, Ronin-Walknowska E. Personality traits assessed by the NEO Five-Factor Inventory (NEO-FFI) as part of the perinatal depression screening program. *Med Sci Monit* 2010; 16: PH77-81.
- Vliegen N, Luyten P, Besser A, Casalin S, Kempke S, Tang E. Stability and change in levels of depression and personality: a follow-up study of postpartum depressed mothers that were hospitalized in a mother-infant unit. *J Nerv Ment Dis* 2010; 198: 45-51.
- Harville E, Xiong X, Buekens P. Disasters and perinatal health: a systematic review. *Obstet Gynecol Surv* 2010; 65: 713-28.
- Sun F, Zhu J, Tao H, Ma Y, Jin W. A systematic review involving 11,187 participants evaluating the impact of COVID-19 on anxiety and depression in pregnant women. *J Psychosom Obstet Gynaecol* 2021; 42: 91-9.
- Bottemanne H, Vahdat B, Jouault C, Tibi R, Joly L. Becoming a mother during COVID-19 pandemic: how to protect maternal mental health against stress factors. *Front Psychiatry* 2022; 12: 764207.
- Wei SQ, Bilodeau-Bertrand M, Liu S, Auger N. The impact of COVID-19 on pregnancy outcomes: a systematic review and meta-analysis. *CMAJ* 2021; 193: E540-E548.
- Schwartz DA, Avvad-Portari E, Babál P, et al. Placental tissue destruction and insufficiency from COVID-19 causes stillbirth and neonatal death from hypoxic-ischemic injury. *Arch Pathol Lab Med* 2022; 146: 660-76.
- Hornig M, Bresnahan MA, Che X, et al. Prenatal fever and autism risk. *Mol Psychiatry* 2018; 23: 759-66.
- Brown AS, Patterson PH. Maternal infection and schizophrenia: implications for prevention. *Schizophr Bull* 2011; 37: 284-90.
- Ayubi E, Mansori K. Maternal infection during pregnancy and attention-deficit hyperactivity disorder in children: a systematic review and meta-analysis. *Iran J Public Health* 2022; 51: 2674-87.
- Faul F, Erdfelder E, Buchner A, Lang AG. Statistical power

- analyses using G*Power 3.1: tests for correlation and regression analyses. *Behav Res Methods* 2009; 41: 1149-60.
20. John OP, Donahue EM, Kentle RL. Big Five Inventory (BFI). Versions 4a and 5. Berkeley, CA: University of California, 1991.
 21. Weiss DS, Marmar CR. The Impact of Event Scale-Revised. In: Wilson JP, Keane TM (eds). *Assessing psychological trauma and PTSD*. New York: The Guilford Press, 1997.
 22. Kroencke L, Geukes K, Utesch T, Kuper N, Back MD. Neuroticism and emotional risk during the COVID-19 pandemic. *J Res Pers* 2020; 89: 104038.
 23. Soto CJ. How replicable are links between personality traits and consequential life outcomes? The Life Outcomes of Personality Replication Project. *Psychol Sci* 2019; 30: 711-27.
 24. Leger KA, Charles ST, Turiano NA, Almeida DM. Personality and stressor-related affect. *J Pers Soc Psychol* 2016; 111: 917-28.
 25. Bellintier JA, Mund M, Wrzus C. The role of extraversion and neuroticism for experiencing stress during the third wave of the COVID-19 pandemic. *Curr Psychol* 2021 Dec 9: 1-11.
 26. Yang Q, Kanjanarat P, Wongpakaran T, et al. Fear of COVID-19 and perceived stress: the mediating roles of neuroticism and perceived social support. *Healthcare (Basel)* 2022; 10: 812.
 27. Campbell SB, Cohn JF, Meyers T. Depression in first-time mothers: mother-infant interaction and depression chronicity. *Dev Psychol* 1995; 31: 349-57.
 28. Gelfand DM, Teti DM. The effects of maternal depression on children. *Clin Psychol Rev* 1990; 10: 329-53.
 29. Priel A, Djalovski A, Zeev-Wolf M, Feldman R. Maternal depression impairs child emotion understanding and executive functions: the role of dysregulated maternal care across the first decade of life. *Emotion* 2020; 20: 1042-58.
 30. van der Waerden J, Bernard JY, De Agostini M, et al. EDEN Mother-Child Cohort Study Group. Persistent maternal depressive symptoms trajectories influence children's IQ: The EDEN mother-child cohort. *Depress Anxiety* 2017; 34: 105-17.
 31. Junge C, Garthus-Niegel S, Slinning K, Polte C, Simonsen TB, Eberhard-Gran M. The impact of perinatal depression on children's social-emotional development: a longitudinal study. *Matern Child Health J* 2017; 21: 607-15.
 32. Hay DF, Pawlby S, Sharp D, Asten P, Mills A, Kumar R. Intellectual problems shown by 11-year-old children whose mothers had postnatal depression. *J Child Psychol Psychiatry* 2001; 42: 871-89.
 33. Pearson J, Tarabulsky GM, Bussièrès EL. Foetal programming and cortisol secretion in early childhood: a meta-analysis of different programming variables. *Infant Behav Dev* 2015; 40: 204-15.
 34. Prenoveau JM, Craske MG, West V, et al. Maternal postnatal depression and anxiety and their association with child emotional negativity and behavior problems at two years. *Dev Psychol* 2017; 53: 50-62.
 35. Whelan YM, Leibenluft E, Stringaris A, Barker ED. Pathways from maternal depressive symptoms to adolescent depressive symptoms: the unique contribution of irritability symptoms. *J Child Psychol Psychiatry* 2015; 56: 1092-100.
 36. Martucci M, Aceti F, Giacchetti N, Scarselli V, Sogos C. A link between parental psychopathology and preschool depression: take care of parents to take care of children. *Children (Basel)* 2023; 10: 150.
 37. Martucci M, Aceti F, Giacchetti N, Sogos C. The mother-baby bond: a systematic review about perinatal depression and child developmental disorders. *Riv Psichiatr* 2021; 56: 223-36.
 38. Lattanzi GM, Provini L, Williams R, Aceti F, Giacchetti N. Personality structure and attachment models of women who kill their children. A systematic review on maternal filicide. *Child Abuse Negl* 2020; 106: 104532.
 39. Giacchetti N, Pancheri C, Mazza C, et al. Differences in MMPI-2 personality profiles among filicide and homicide women. *Riv Psichiatr* 2020; 55: 152-60.
 40. Martucci M, Fava G, Giacchetti N, et al. Perinatal depression as a risk factor for child developmental disorders: a cross-sectional study. *Riv Psichiatr* 2021; 56: 321-7.
 41. Giangiacomo E, Visaggi MC, Aceti F, et al. Early neuropsychomotor therapy intervention for theory of mind and emotion recognition in neurodevelopmental disorders: a pilot study. *Children (Basel)* 2022; 9: 1142.
 42. Rocchi G, Serio V, Carluccio GM, et al. La regolazione epigenetica della relazione primaria. *Riv Psichiatr* 2015; 50: 155-60.
 43. Vitali M, Tedeschini E, Mistretta M, et al. Adjunctive pregabalin in partial responders with major depressive disorder and residual anxiety. *J Clin Psychopharmacol* 2013; 33: 95-8.

Corresponding author:
Melania Martucci
Department of Human Neurosciences
Sapienza University of Rome
Viale dell'Università 30
00185 Rome, Italy
E-mail: melania.martucci@uniroma1.it