

Diagnosis and management of tardive dyskinesia: from research to clinical practice

ANDREA FAGIOLINI¹, PAOLO BARONE², ANTONELLO BELLOMO³, EMI BONDI⁴, CARLO COLOSIMO⁵, GIOVANNI FABBRINI⁶, GIUSEPPE MAINA⁷, MAURIZIO POMPILI⁸, MICHELE TINAZZI⁹, ANTONIO VITA¹⁰, ALESSANDRO CUOMO¹

¹University of Siena, Italy; ²Department of Neurology, University of Salerno, Italy; ³Department of Clinical and Experimental Medicine, University of Foggia, Italy; ⁴Department of Mental Health, ASST Bergamo, Italy; ⁵Department of Neurology, Santa Maria Hospital, Terni, Italy; ⁶Department Human Neurosciences, Sapienza University of Rome and IRCCS Neuromed; ⁷Psychiatry Division, University hospital San Luigi Gonzaga of Orbassano - School of Medicine, University of Turin, Italy; ⁸Department of Neurosciences, Mental Health and Sensory Organs, Suicide Prevention Centre, Sant'Andrea Hospital, Sapienza University of Rome, Italy; ⁹Department of Neurology, AO University of Verona, Italy; ¹⁰University of Brescia, ASST Spedali Civili of Brescia, Italy.

Summary. Tardive dyskinesia (TD) is a chronic, often disabling hyperkinetic movement disorder associated with prolonged use of dopamine receptor blocking agents (DRBAs), particularly antipsychotics (APs) for psychiatric disorders such as schizophrenia and bipolar disorder. It manifests as abnormal, involuntary movements, often involving the orofacial region, extremities, or trunk, and is associated with significant physical and psychosocial impairment. TD is primarily linked to dopamine receptor hypersensitivity, oxidative stress, and genetic susceptibility, with a higher prevalence in patients treated with first-generation APs. However, second-generation APs (SGAs) have not eliminated the risk entirely, particularly in older adults and those with prolonged exposure. Diagnosis relies on clinical assessments such as the Abnormal Involuntary Movement Scale (AIMS) and comprehensive neurological evaluations. Treatment guidelines emphasize early detection, prevention through minimal effective doses of APs, and the use of VMAT2 inhibitors (vesicular monoamine transporter 2 inhibitors) as a first-line therapy in moderate-to-severe cases. VMAT2 inhibitors reduce dopamine signaling dysregulation without directly blocking D2 receptors, effectively managing symptoms in many patients. For treatment-resistant cases, deep brain stimulation and other non-pharmacological interventions offer promising alternatives. Current research underscores the complexity of TD's pathophysiology and the need for personalized approaches. Future directions include developing biomarkers for risk stratification, refining therapeutic strategies, and optimizing long-term outcomes through multidisciplinary care.

Key words. Antipsychotic-induced movement disorders, dopamine receptor blocking agents, hyperkinetic movement disorders, tardive dyskinesia, VMAT2 inhibitors.

Introduction

Tardive dyskinesia is a persistent and often disabling hyperkinetic movement disorder associated

Diagnosi e gestione della discinesia tardiva: dalla ricerca alla pratica clinica.

Riassunto. La discinesia tardiva (TD) è un disturbo del movimento ipercinetico cronico e spesso invalidante, associato all'uso prolungato di agenti bloccanti i recettori della dopamina (DRBA), principalmente antipsicotici (AP) utilizzati per trattare schizofrenia e disturbo bipolare. Si manifesta con movimenti involontari anomali, spesso localizzati nella regione orofacciale, alle estremità o al tronco, e comporta un impatto significativo sulla salute fisica e psicosociale. La TD è legata principalmente all'ipersensibilità dei recettori dopaminergici, allo stress ossidativo e a fattori genetici, con una prevalenza maggiore nei pazienti trattati con antipsicotici di prima generazione. Tuttavia, anche gli antipsicotici di seconda generazione (SGA) non hanno eliminato completamente il rischio, soprattutto nei pazienti anziani e in trattamenti prolungati. La diagnosi si basa su valutazioni cliniche, come la scala Abnormal Involuntary Movement Scale (AIMS), e su un esame neurologico approfondito. Le linee guida terapeutiche si concentrano sulla prevenzione tramite dosi minime efficaci di AP e sull'uso precoce di inibitori VMAT2 (inibitori del trasportatore vescicolare delle monoamine di tipo 2), considerati il trattamento di prima linea nei casi moderati e severi. Gli inibitori VMAT2 modulano la disregolazione della segnalazione dopaminergica senza bloccare direttamente i recettori D2, offrendo una gestione efficace dei sintomi nella maggior parte dei pazienti. Nei casi resistenti al trattamento, la stimolazione cerebrale profonda e altre opzioni non farmacologiche rappresentano alternative promettenti. Le ricerche attuali evidenziano la complessità della patofisiologia della TD e la necessità di approcci personalizzati. Le prospettive future includono lo sviluppo di biomarcatori per la stratificazione del rischio, il miglioramento delle strategie terapeutiche e l'ottimizzazione degli esiti a lungo termine attraverso cure multidisciplinari.

Parole chiave. Agenti bloccanti i recettori della dopamina, discinesia tardiva, disturbi del movimento indotti da antipsicotici, disturbi del movimento ipercinetico, inibitori VMAT2.

with prolonged exposure to dopamine receptor blockers, particularly antipsychotics (APs) used to treat psychiatric disorders such as schizophrenia and bipolar disorder. Research suggests that second-generation APs have a lower risk of tardive dyskinesia

than first-generation APs¹. Tardive syndromes can manifest in a variety of ways, including oro-buccal-lingual stereotypy, akathisia, dystonia, tics, tremor, and chorea². Tardive dyskinesia (TD) is the most common tardive syndrome and is characterized by abnormal involuntary movements, typically manifesting as involuntary continuous orofacial movements and movements of the extremities or trunk³. It is a serious, potentially permanent neurological hyperkinetic movement disorder that occurs after months or years of psychotropic medication use⁴. Despite the expectation that atypical APs would reduce the prevalence of TD, data show that TD remains a significant challenge due to the persistent nature of its symptoms and resistance to various treatment modalities, including AP discontinuation⁵. It is important to note that TD can occur independently of AP use: in the pre-AP era, spontaneous dyskinesia was observed in patients with psychosis⁶ and prolonged use of metoclopramide and prochlorperazine can also lead to TD due to their dopamine antagonist properties⁷.

In addition, TD often transiently worsens after cessation of the causative DRBA⁸. The etiopathogenesis of TD remains unclear, although it is associated with long-term AP use⁹. While the new generation of atypical AP agents may potentially eliminate concerns about TD in the future, it currently remains a significant clinical problem for patients, their caregivers, and physicians¹⁰. The effects of antipsychotics are inherently multidimensional, encompassing diverse neurobiological targets that extend beyond dopamine receptor modulation. These include immunomodulatory actions, particularly on microglia, as well as interactions with neuroactive steroids and inflammatory pathways, which may contribute to symptom improvement in schizophrenia spectrum disorders^{11,12}. Effective management of TD is critical, as variants such as tardive dystonia and tardive akathisia tend to be more severe and challenging to treat than typical TD¹³. Tardive dystonia consists of sustained muscle contractions and irregular postures, whereas tardive akathisia consists of motor restlessness and subjective discomfort. AP drug withdrawal rarely benefits patients with these variants¹³, but reserpine, first-generation VMAT2 inhibitors, β -blockers, benzodiazepines, or clozapine may benefit certain patients¹⁴. Patients with focal tardive dystonia may also improve with local injection of botulinum neurotoxin.

It is also essential to differentiate TD from other movement disorders. The differential diagnosis of TD is based on several factors, including a history of exposure to dopamine receptor blocking agents (DRBAs), the duration of such exposure, the specific characteristics of the movement abnormalities, and a history of related systemic or neurologic conditions^{2,8}.

In summary, TD is a complex and challenging movement disorder associated with long-term use of AP medications. While advances have been made in the understanding and managing TD, further research is needed to improve treatment outcomes and quality of life for individuals affected by this condition.

Materials and methods

This narrative review synthesizes current evidence on TD, focusing on its diagnosis, pathophysiology, and management strategies. The authors conducted a comprehensive literature search using PubMed, Scopus, and Web of Science databases for articles published through September 2024. The search strategy combined terms related to “tardive dyskinesia”, “VMAT2 inhibitors”, “dopamine receptor blockers”, “antipsychotic-induced movement disorders”, and their variants. We included peer-reviewed studies, clinical trials, systematic reviews, and meta-analyses that addressed TD’s epidemiology, etiology, clinical features, and treatment options. Priority was given to publications from the past decade, although seminal older papers were included when relevant. The search was limited to English-language publications. We focused particularly on high-quality evidence, including randomized controlled trials, meta-analyses, and current clinical practice guidelines. Special attention was paid to studies examining the efficacy and safety of VMAT2 inhibitors, as these represent the newest approved treatments for TD. The narrative synthesis of the literature aims to provide clinicians with practical, evidence-based guidance for the diagnosis and management of TD.

Overview of the paper

This article provides a comprehensive review of the current understanding of TD, a persistent and often disabling movement disorder associated with prolonged exposure to dopamine receptor blockers, particularly AP drugs¹⁵. We discuss the epidemiology, risk factors, pathophysiology, clinical features, diagnostic criteria and tools, and treatment of TD^{16,17}. The burden of TD on patients and healthcare systems is reviewed, highlighting its impact on quality of life, social stigma, and economic costs^{18,19}. Current guidelines for managing TD are reviewed, focusing on prevention strategies, dose reduction or discontinuation of causative agents, and the use of VMAT2 inhibitors^{20,21}. Recent advances in understanding the pathophysiology of TD are also reviewed, including the role of dopamine receptor hypersensitivity, GABA insufficiency, oxidative stress, and neuronal toxicity²²⁻²⁶. We used a narrative approach for this review.

The literature search was conducted using specific keywords such as “tardive dyskinesia”, “VMAT2 inhibitors”, “dopamine receptor blockers”, and “antipsychotic-induced movement disorders”. We searched databases, including PubMed, Scopus, and Web of Science, for studies published through September 2024. The search aimed to identify relevant clinical trials, reviews, and meta-analyses focusing on tardive dyskinesia’s epidemiology, pathophysiology, and management. Although this review is not systematic, we have carefully selected the most relevant literature to support our discussion and expert opinion. We conclude with expert opinions on future directions for research and treatment, emphasizing the need for improved prevention, early detection, and individualized management of TD in clinical practice.

Epidemiology

The risk of developing TD differs between first-generation APs (FGAs) and second-generation APs (SGAs)²⁷. Multiple studies and meta-analyses have consistently shown that SGAs are associated with a lower risk of TD compared to FGAs, with a cumulative annual incidence of 5.4%-7.7% with FGAs compared to 0.8%-3.0% with SGAs²⁷. A meta-analysis by Carbon et al. (2017) reported that global mean TD prevalence was 25.3% across all studies, with lower rates in patients treated with SGAs (20.7%) compared to those treated with FGAs (30.0%)^{27,28}. The reduced risk of TD with SGAs is related to their lower affinity for dopamine D2 receptors and their effects on other neurotransmitter systems, such as serotonin^{29,30}. However, it is important to note that the advantage of SGAs over FGAs in terms of TD risk may not be as substantial as initially believed, and TD can still occur with SGAs, particularly in vulnerable populations such as older adults and those with a longer duration of AP exposure^{22,30,31}. In addition, some studies have suggested that the risk of TD may vary between SGAs, with risperidone and olanzapine possibly having a lower risk than other SGAs^{29,32,33}. Other antipsychotics, such as aripiprazole, brexpiprazole, cariprazine, and lurasidone, are less commonly associated with TD, although they may predispose patients to other side effects, such as akathisia.

Sex differences in the risk of TD have been reported in some studies, with a higher prevalence in women compared with men, especially in older age groups^{34,35}. A meta-analysis by Yassa and Jeste³⁴ found that the prevalence of TD was 26.6% in women and 21.6% in men, with the sex difference being more pronounced in patients over 50 years of age. However, other studies have not found significant sex differences in the risk of TD⁵⁷. Older age is a well-established risk factor for TD, with higher prevalence and

incidence observed in older patients^{34,36}. The risk of TD appears to increase significantly after age 50, with some studies suggesting a cut-off age of 55 years^{34,36}. A prospective study by Jeste et al.³⁶ found that the incidence of TD was 25% in patients aged 55 years and older compared to 10% in younger patients.

Despite the reduced risk with SGAs, TD remains a significant concern in patients requiring long-term AP treatment, and regular monitoring for the onset of TD symptoms is critical for early detection and management³¹.

Etiology and pathophysiology

The etiology of TD is multifactorial, involving a complex interplay of genetic predisposition, oxidative stress, and neurochemical imbalances, particularly in the dopaminergic system³⁷. Chronic blockade of dopamine receptors, particularly D2 receptors, by dopamine receptor blockers such as APs is thought to lead to compensatory upregulation and hypersensitivity of these receptors, contributing to TD^{32,33}. Genetic factors, including polymorphisms in genes related to dopamine receptors, drug metabolism, and oxidative stress, may influence an individual’s susceptibility to developing TD²¹. Candidate gene studies in TD have identified a number of promising leads, including CYP2D6, DRD2, SLC18A2, HTR2A, and HSPG2. Other genes, particularly in the γ -aminobutyric acid (GABA) and glutamate systems, may be worthy of additional interrogation, especially given the observed co-transmission of dopamine and GABA through vesicular co-packaging by VMAT2 in striatal dopaminergic neurons³⁷.

Oxidative stress, resulting from an imbalance between the production of reactive oxygen species and the body’s antioxidant defenses, has been implicated in the pathogenesis of TD³⁸. Chronic exposure to DRBAs may increase dopamine turnover and oxidative stress in the basal ganglia, resulting in neurotoxicity and neuroinflammation^{38,39}. These pathophysiological mechanisms, along with age-related changes in the brain and other environmental factors, may collectively contribute to the development of TD in susceptible individuals^{32,33}.

Clinical features

The symptoms of TD (figure 1) can vary in severity and significantly impact the quality of life of affected individuals. Common manifestations include repetitive, involuntary movements such as facial grimacing, tongue protrusion, lip smacking, and rapid eye blinking^{2,40}. These abnormal movements predominantly affect the orofacial region but

can also extend to involve the trunk and extremities, leading to more generalized motor impairment^{2,40}. Individuals with TD may encounter challenges in daily activities such as eating, speaking and walking. They may also face social stigma and isolation due to the visible nature of the symptoms⁴¹. The presentation of TD symptoms can range from mild and intermittent movements to severe, continuous, and disabling manifestations^{2,40}. Illustrative cases demonstrate a spectrum of TD manifestations, from isolated perioral movements to generalized chorea and dystonia affecting multiple body regions^{2,42}. Oro-bucco-lingual dyskinesia, often considered the classic form of tardive dyskinesia, was observed in 72% of cases². Tardive tremor affected 30% of the patients, while 22% experienced tardive akathisia, and 16% suffered from tardive dystonia². Less common manifestations included tardive tics and myoclonus, affecting 4% and 1% of patients, respectively². Additionally, 35% of individuals presented with a combination of two or more tardive syndromes². The impact of TD on daily functioning and quality of life is substantial, with patients reporting physical discomfort, emotional distress, and limitations in social and occupational engagement⁴¹.

Diagnosis

The diagnosis of TD (figure 2) relies primarily on a comprehensive clinical assessment, which includes a detailed history of exposure to dopamine receptor-blocking agents and a thorough neurological examination to identify characteristic involuntary movements^{2,4,43}. The Abnormal Involuntary Movement Scale (AIMS) (Appendix 1) is a validated tool commonly used (particularly in clinical research) to assess the severity of TD symptoms, with scores ranging from 0 (no dyskinesia) to 4 (severe dyskinesia) across 12 items evaluating involuntary movements in various body regions (including face,

extremities, and trunk), as well as overall severity, impact and awareness of abnormal movements^{44,45}. Items 1 through 7 quantify abnormal movements involving the face, extremities, and trunk⁴⁵. An AIMS score ≥ 2 in 2 or more body areas or a score of 3 or 4 in at least 1 body region indicates an abnormal hyperkinetic movement like TD, according to Schooler-Kane criteria⁴⁵. Additional diagnostic tools such as the Extrapyrarnidal Symptom Rating Scale (ESRS) and the Dyskinesia Identification System: Condensed User Scale (DISCUS) offer further insights into TD severity and its impact on daily functioning^{46,47}. Clinical history and physical examination play a crucial role in confirming the diagnosis of TD and differentiating it from other movement disorders that may present similarly, such as Huntington's disease, primary dystonia or Wilson's disease^{2,4}. A thorough differential diagnosis is essential to ensure appropriate management and treatment of the underlying condition⁴⁸ (Appendix 2).

Burden of disease

The impact of TD on patients is substantial, affecting their physical, emotional, and social well-being. The involuntary movements associated with TD can lead to stigmatization, social withdrawal, and a decline in quality of life^{48,49}. Individuals with TD often experience embarrassment, social isolation, and difficulties in daily activities such as eating, speaking, and walking^{48,49}. The psychological burden of TD is significant, with patients reporting higher rates of depression, anxiety, and low self-esteem^{1,48}. Furthermore, the economic consequences of TD include increased healthcare costs due to ongoing management and potential disability⁴⁹. Patients with TD demonstrate higher rates of healthcare resource utilization, including more frequent visits to healthcare providers, increased use of emergency services,




Figure 1. Symptoms of Tardive Dyskinesia (TD).




Figure 2. Diagnosis of Tardive Dyskinesia (TD).

Legend: DRBAs= dopamine receptor blocking agents; AIMS= Abnormal Involuntary Movement Scale; ESRS= Extrapyramidal Symptom Rating Scale; DISCUS= Dyskinesia Identification System: Condensed User Scale.

and longer hospital stays compared to those without TD^{41,49}. TD's social and economic impacts extend beyond the individual patient, affecting their families and the healthcare system as a whole^{41,49}.

Guidelines

The management of TD is directed by the latest clinical guidelines that emphasize early detection and the use of the lowest effective dose of AP medications. The American Academy of Neurology (AAN) and the American Psychiatric Association (APA) provide recommendations for assessing and treating TD, advocating for regular screening and considering VMAT2 inhibitors as treatment options^{45,48}. The AAN guidelines underscore the importance of monitoring for TD signs and suggest the use of VMAT2 inhibitors when appropriate^{45,48}. APA guidelines recommend that all patients receiving AP medications should be regularly monitored for movement disorders^{45,48}. The APA recommends that patients treated with FGAs be monitored every 6 months, while patients on SGA drugs should be evaluated every 12 months^{45,48}. According to the APA guidelines, a closer monitoring for abnormal movements should be performed for patients at increased risk for TD (e.g., older adults, presence of affective disorder) receiving AP medications^{45,48}. These high-risk patients should be assessed for movement disorders every 3 months if taking FGAs or every 6 months if receiving SGAs^{45,48}.

In terms of treatment strategies, APA recommends VMAT2 inhibitors for patients with moderate to severe or disabling TD associated with AP therapy^{45,48}. Both guidelines stress the significance of early detection and minimal effective AP doses to reduce the risk of TD development^{45,48}. Best practices for monitoring include regular clinical assessments and standardized rating scales such as the AIMS^{45,48}. Early intervention strategies, such as AP dose reduction, switching to lower-risk APs, or initiating VMAT2 inhibitors, are crucial in preventing or alleviating the long-term consequences of TD^{45,48}. Indeed, switching from an FGA, particularly haloperidol, to an SGA with a lower D2 affinity, such as clozapine or quetiapine, may reduce TD symptoms⁴⁸.

Regarding VMAT2 inhibitors, there is good evidence to support a favourable benefit-risk profile for novel VMAT2 inhibitors as a treatment option for TD⁴⁸. Novel VMAT2 inhibitors should be considered as first-line treatment for TD⁴⁸. Instead, there is limited evidence for the use of first-generation VMAT2 inhibitors for the treatment of TD⁴⁸. The current evidence suggests that, while first-generation VMAT2 inhibitors may be helpful for TD, their use is associated with more adverse effects than novel VMAT2 inhibitors⁴⁸.

An interdisciplinary approach involving collaboration among psychiatrists, neurologists, and primary care providers is essential for the optimal management of TD and the underlying psychiatric condition.

Treatment

The treatment of TD involves various approaches and considerations. One approach to managing TD is to discontinue AP treatment or reduce its dosage. However, these options may not always be feasible due to the potential risk of exacerbation of symptoms or negative impact on psychiatric status⁵⁰. Recommendations for treating TD, including the use of adrenergic uptake inhibitors and AP agents, are provided in systematic reviews⁴⁸. Introducing new treatments, such as VMAT2 inhibitors, has revitalized interest in recognizing and managing TD^{45,48}. The pharmacological approach to treating TD has explored various medications and interventions to alleviate symptoms and improve patient outcomes. VMAT2 inhibitors have shown promise in managing TD, with studies highlighting their efficacy and safety profiles^{45,48,51}.

Additionally, the use of botulinum toxin injections has been considered in the treatment of TD, particularly in cases of focal dyskinesia². The botulinum toxin acts at the alpha motoneuron nerve terminals at the injected site exerting its effect as local muscle relaxant⁵². The efficacy and tolerability of VMAT2 inhibitors for TD have been systematically reviewed, shedding light on their potential as a treatment option^{45,48}. VMAT2 inhibition works by reducing dopamine stimulation without directly blocking D2 receptors. This mechanism decreases the overstimulation of D2 receptors in the indirect pathway, which in turn reduces inhibition of the “stop” signal.

Additionally, VMAT2 inhibition benefits in the direct pathway by amplifying “go” signals, as dopamine’s effects on D1 receptors are enhanced⁵³. Furthermore, vitamin E’s role in treating TD has been investigated, with meta-analyses aiming to determine its effectiveness in alleviating symptoms⁵⁴. Vitamin E may help reduce the damage caused by the overproduction of cytotoxic free radicals and may prevent or lessen the severity of TD, especially in individuals who have experienced the onset of the condition within the past five years⁵⁴. The use of anticholinergic medications should be discouraged, and it may worsen TD. Moreover, case studies suggest that neurostimulation via deep brain stimulation (DBS) or repetitive transcranial magnetic stimulation (rTMS) may be an effective option in severe, refractory cases of TD⁵⁵. An interdisciplinary approach involving psychiatrists, neurologists, and primary care providers is crucial for the optimal management of TD and for selecting appropriate pharmacological interventions tailored to individual patient needs.

Expert opinion and conclusions

In summary, tardive dyskinesia (TD) remains a challenging disease to manage, with significant impli-

cations for patients, their caregivers, and healthcare systems. Advances in understanding the pathophysiology of TD, including the role of dopamine receptor hypersensitivity, GABA insufficiency, and oxidative stress, have led to targeted treatments such as VMAT2 inhibitors. VMAT2 inhibitors offer hope for better management of TD symptoms and improved patient outcomes. However, challenges remain in optimizing TD prevention, early detection, and individualized treatment in clinical practice. Future research should focus on identifying predictive biomarkers of TD susceptibility and treatment response, comparing the long-term efficacy and safety of VMAT2 inhibitors, and exploring novel therapeutic approaches such as deep brain stimulation and transcranial magnetic stimulation. Expert evidence highlights the importance of regular screening for TD (through regular use of the AIMS scale in clinical practice, as it is a simple, rapid and reliable tool for diagnosis and monitoring of TD), using the lowest effective dose of APs, and considering VMAT2 inhibitors for patients with moderate to severe TD. Recommendations for clinical practice include adopting a prevention-oriented approach, educating patients and caregivers about the risk of TD, and collaborating with movement disorder specialists for complex cases. In terms of managing TD in current clinical practice, experts suggest a stepwise approach^{56,57}. First, prevention strategies should be employed, such as using the lowest effective dose of antipsychotics and regularly monitoring for early signs of TD using the AIMS scale⁵⁶. If TD is detected, the offending antipsychotic should be discontinued or switched to a lower-risk alternative, such as clozapine or quetiapine, if clinically feasible⁵⁷. For patients with moderate to severe TD, VMAT2 inhibitors should be considered as first-line pharmacological treatment⁵⁷. Novel VMAT2 inhibitors have demonstrated efficacy in reducing TD severity and are generally well-tolerated^{51,58}. In cases where TD is refractory to pharmacological interventions, non-pharmacological options such as deep brain stimulation may be explored in collaboration with movement disorder specialists⁵⁹. Management of tardive TD requires routine assessment of its impact on daily functioning and quality of life using tools like the Impact of TD Scale. An integrative, interdisciplinary approach – combining psychiatrists, neurologists, and primary care providers – should address motor, psychological, and social aspects of TD. Incorporating the patient’s perspective into treatment decisions fosters individualized care. Prevention and early detection, alongside personalized pharmacological and non-pharmacological strategies, are critical for minimizing TD’s burden and improving outcomes. Alongside pharmacological management, it is crucial to educate and support patients and their caregivers⁵⁶. This includes informing them about the potential risk of TD with long-term antipsychotic use, the impor-

Recommendations for assessing TD impact in clinical practice.

Routinely assess the impact of TD on patients' daily functioning and quality of life as part of standard care, in addition to assessing TD severity.

Use a standardized, easy-to-administer tool such as the Impact of TD Scale to systematically assess the frequency and distress associated with specific functional consequences of TD symptoms.

Key domains to assess include social, physical, psychological, and cognitive functioning, as well as activities of daily living.

Incorporate the patient's perspective on the impact of TD into treatment, decision-setting, decision-making, and goal-setting.

Managing TD requires an integrative approach with routine impact assessments, interdisciplinary care, patient-centered treatment, and prevention strategies to minimize its burden and improve outcomes.

Collaborate with interdisciplinary teams, including psychiatrists, neurologists, and primary care providers, to optimize TD management and address the multifaceted impact of TD on patients' lives.

Consider alternative analyses of Abnormal Involuntary Movement Scale (AIMS) data, such as effect size and minimal clinically significant difference, to improve TD assessment's clinical relevance and interpretability in practice.

tance of regular monitoring, and the available treatment options. Patients should be encouraged to report any involuntary movements to their healthcare providers promptly⁵⁴. In summary, the management of TD in current clinical practice involves a combination of prevention, early detection, individualized pharmacological and non-pharmacological interventions, and patient and caregiver education. By adopting these strategies, clinicians can work toward minimizing the burden of TD and improving the quality of life for patients with this challenging condition.

Funding: this work received an unrestricted grant from Teva Italia. The funders had no role in study design, data collection and analysis, decision to publish, or manuscript preparation.

Authorship: all named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship for this article, take responsibility for the integrity of the work as a whole, and have given their approval for this version to be published. All authors contributed equally to the manuscript.

Conflict of interests: the authors wish to disclose that in the last five years they have received lectures and advisory board honoraria or have engaged in clinical trial activities with the following companies, all of which are unrelated to this article. AB: Angelini, Janssen, Lundbeck, Rovi, Viatri, Teva; EB: Fidia, Janssen, Lundbeck, Rovi, Teva, Viatri; CC: BIAL, Ibsen, Teva; AC: Angelini, GlaxoSmithKline, Italfarmaco, Lundbeck, Janssen, Mylan, Neuraxpharm, Otsuka, Pfizer, Recordati, Sanofi Aventis, Viatri; AF: Angelini, Aspen, Biogen, Boehringer Ingelheim, Janssen, Lundbeck, Neuraxpharm, Otsuka, Pfizer, Recordati, Viatri; GM: Angelini, Lundbeck,

Janssen, Otsuka, Boehringer Ingelheim, Biohaven Pharma, Teva, Rovi, Italfarmaco, Viatri; MP: Angelini Pharma, Allergan, Janssen, Lundbeck, Merck Sharp & Dohme, Otsuka, Rovi, Pfizer, Fidia, Viatri, Recordati, Boehringer Ingelheim, Newron and GSK; MT: Zambon, Lusofarmaco, AbbVie, Bial, Chiesi; AV: Alkermes, Angelini, Boehringer Ingelheim, Janssen-Cilag, Lundbeck, Otsuka, Roche, Rovi Pharma, Teva. PB and GF have no conflict of interests to declare.

Editorial and other assistance: the authors want to thank Sanitanova for editorial assistance.

References

- Correll CU, Leucht S, Kane JM. Lower risk for tardive dyskinesia associated with second-generation antipsychotics: a systematic review of 1-year studies. *AJP* 2004; 161: 414-25.
- Waln O, Jankovic J. An update on tardive dyskinesia: from phenomenology to treatment. *TOHM* 2013; 3: 03.
- Stegmayer K, Walther S, Van Harten P. Tardive dyskinesia associated with atypical antipsychotics: prevalence, mechanisms and management strategies. *CNS Drugs* 2018; 32: 135-47.
- Khouzam HR. Identification and management of tardive dyskinesia: a case series and literature review. *PMJ* 2015; 127: 726-37.
- Meyer JM. Forgotten but not gone: new developments in the understanding and treatment of tardive dyskinesia. *CNS Spectr* 2016; 21: 13-24.
- Haddad PM, Dursun SM. Neurological complications of psychiatric drugs: clinical features and management. *Hum Psychopharmacol Clin Exp* 2008; 23: S15-S26.
- Cornett EM, Novitch M, Kaye AD, et al. Medication-induced tardive dyskinesia: a review and update. *Ochsner J* 2017; 17: 162-74.
- Perju-Dumbrava L, Kempster P. Movement disorders in psychiatric patients. *BMJ Neurol Open* 2020; 2: e000057.
- Lohr JB, Kuczenski R, Niculescu AB. Oxidative mechanisms and tardive dyskinesia. *CNS Drugs* 2003; 17: 47-62.
- Egan MF, Apud J, Wyatt RJ. Treatment of tardive dyskinesia. *Schizophr Bull* 1997; 23: 583-609.
- Orsolini L, De Berardis D, Volpe U. Up-to-date expert opinion on the safety of recently developed antipsychotics. *Expert Opin Drug Saf* 2020; 19: 981-98.
- Amerio A, Magnani L, Arduino G, et al. Immunomodulatory effects of clozapine: more than just a side effect in schizophrenia. *Curr Neuropharmacol* 2024; 22: 1233-47.
- Gardos G. Managing antipsychotic-induced tardive dyskinesia. *Drug Saf* 1999; 20: 187-93.
- Skidmore F, Weiner WJ, Burke R. Neuroleptic-induced tardive dyskinesia variants. In: Factor SA, Lang AE, Weiner WJ (eds). *Drug induced movement disorders*. Hoboken, NJ: Wiley, 2005.
- Marshall DL, Hazlet TK, Gardner JS, et al. Neuroleptic drug exposure and incidence of tardive dyskinesia: a records-based case-control study. *JMCP* 2002; 8: 259-65.
- Rissardo JP, Vora N, Mathew B, et al. Overview of movement disorders secondary to drugs. *Clin Pract* 2023; 13: 959-76.
- Trottenberg T. Pallidal and thalamic neurostimulation in severe tardive dystonia. *J Neurol Neurosurg Psychiatry* 2001; 70: 557-9.
- Albayrak Y, Hashimoto K. Beneficial effects of sigma-1 agonist fluvoxamine for tardive dyskinesia and tardive akathisia in patients with schizophrenia: report of three cases. *Psychiatry Investig* 2013; 10: 417.
- Elkashef AM, Wyatt RJ. Tardive dyskinesia: possible involvement of free radicals and treatment with vitamin E. *Schizophr Bull* 1999; 25: 731-40.

20. Ghosn O, Ye E, Huege S. Evaluating and managing tardive dyskinesia in the older adult. *Curr Geri Rep* 2021; 10: 108-15.
21. Vijayakumar D, Jankovic J. Drug-induced dyskinesia, Part 2: Treatment of tardive dyskinesia. *Drugs* 2016; 76: 779-87.
22. Stahl SM. Neuronal traffic signals in tardive dyskinesia: not enough "stop" in the motor striatum. *CNS Spectr* 2017; 22: 427-34.
23. Blin J, Baron JC, Cambon H, et al. Striatal dopamine D2 receptors in tardive dyskinesia: PET study. *J Neurol Neurosurg Psychiatry* 1989; 52: 1248-52.
24. Solbrig MV, Koob GF, Lipkin WI. Orofacial dyskinesias and dystonia in rats infected with Borna disease virus; a model for tardive dyskinetic syndromes. *Mol Psychiatry* 1999; 4: 310-2.
25. Chouinard G, Samaha A-N, Chouinard V-A, et al. Antipsychotic-induced dopamine supersensitivity psychosis: pharmacology, criteria, and therapy. *Psychother Psychosom* 2017; 86: 189-219.
26. Tsai G, Goff DC, Chang RW, et al. Markers of glutamatergic neurotransmission and oxidative stress associated with tardive dyskinesia. *AJP* 1998; 155: 1207-13.
27. Solmi M, Pigato G, Kane JM, et al. Treatment of tardive dyskinesia with VMAT-2 inhibitors: a systematic review and meta-analysis of randomized controlled trials. *DDDT* 2018; 12: 1215-38.
28. Carbon M, Hsieh C-H, Kane JM, et al. Tardive dyskinesia prevalence in the period of second-generation antipsychotic use: a meta-analysis. *J Clin Psychiatry* 2017; 78: e264-e278.
29. Caroff SN, Aggarwal S, Yonan C. Treatment of tardive dyskinesia with tetrabenazine or valbenazine: a systematic review. *J Comp Eff Res* 2018; 7: 135-48.
30. Marsden CD, Jenner P. The pathophysiology of extrapyramidal side-effects of neuroleptic drugs. *Psychol Med* 1980; 10: 55-72.
31. Thanvi B, Lo N, Robinson T. Levodopa-induced dyskinesia in Parkinson's disease: clinical features, pathogenesis, prevention and treatment. *Postgrad Med J* 2007; 83: 384-8.
32. Teo JT, Edwards MJ, Bhatia K. Tardive dyskinesia is caused by maladaptive synaptic plasticity: a hypothesis. *Mov Disord* 2012; 27: 1205-15.
33. Al-Saffar A, Lennernäs H, Hellström PM. Gastroparesis, metoclopramide, and tardive dyskinesia: Risk revisited. *Neurogastroenterology Motil* 2019; 31: e13617.
34. Yassa R, Jeste DV. Gender differences in tardive dyskinesia: a critical review of the literature. *Schizophr Bull* 1992; 18: 701-15.
35. Woerner MG, Alvir MJ, Saltz BL, Lieberman JA, Kane JM. Prospective study of tardive dyskinesia in the elderly: rates and risk factors. *Am J Psychiatry* 1998; 155: 1521-8.
36. Jeste DV, Caligiuri MP, Paulsen JS, et al. Risk of tardive dyskinesia in older patients: a prospective longitudinal study of 266 outpatients. *Arch Gen Psychiatry* 1995; 52: 756-65.
37. Zai CC, Maes MS, Tiwari AK, et al. Genetics of tardive dyskinesia: promising leads and ways forward. *J Neurol Sci* 2018; 389: 28-34.
38. Margolese HC, Chouinard G, Kolivakis TT, et al. Tardive Dyskinesia in the era of typical and atypical antipsychotics. Part 1: Pathophysiology and mechanisms of induction. *Can J Psychiatry* 2005; 50: 541-7.
39. Montastruc J, Llau M, Rascol O, et al. Drug-induced parkinsonism: a review. *Fundam Clin Pharmacol* 1994; 8: 293-306.
40. O'Brien CF, Jimenez R, Hauser RA, et al. NBI-98854, a selective monoamine transport inhibitor for the treatment of tardive dyskinesia: a randomized, double-blind, placebo-controlled study. *Mov Disord* 2015; 30: 1681-7.
41. Peckham AM, Nicewonder JA. VMAT2 inhibitors for tardive dyskinesia - Practice implications. *J Pharm Pract* 2019; 32: 450-7.
42. Haji Rahman R, Dharmapuri S. Oral dyskinesia in a pediatric patient following concurrent use of neuroleptics and stimulants: treatment strategy considerations to avert avoidable adverse side effects. *Cureus* 2023; 15: e38294.
43. Kay SR, Fiszbein A, Opler LA. The Positive and Negative Syndrome Scale (PANSS) for schizophrenia. *Schizophr Bull* 1987; 13: 261-76.
44. Lohr JB, Eidt CA, Abdulrazzaq Alfaraj A, et al. The clinical challenges of akathisia. *CNS Spectr* 2015; 20: 1-16.
45. Correll CU, Kane JM, Citrome LL. Epidemiology, prevention, and assessment of tardive dyskinesia and advances in treatment. *J Clin Psychiatry* 2017; 78: 1136-47.
46. Ward KM, Citrome L. Antipsychotic-related movement disorders: drug-induced parkinsonism vs. tardive dyskinesia - Key differences in pathophysiology and clinical management. *Neurol Ther* 2018; 7: 233-48.
47. Macaluso M, Flynn A, Preskorn S. Determining whether a definitive causal relationship exists between aripiprazole and tardive dyskinesia and/or dystonia in patients with major depressive disorder, Part 4: Case Report Data. *J Psychiatr Pract* 2016; 22: 203-20.
48. Ricciardi L, Pringsheim T, Barnes TRE, et al. Treatment recommendations for tardive dyskinesia. *Can J Psychiatry* 2019; 64: 388-99.
49. Browne S, Roe M, Lane A, et al. Quality of life in schizophrenia: relationship to sociodemographic factors, symptomatology and tardive dyskinesia. *Acta Psychiatr Scand* 1996; 94: 118-24.
50. Hauser RA, Factor SA, Marder SR, et al. KINECT 3: a phase 3 randomized, double-blind, placebo-controlled trial of valbenazine for tardive dyskinesia. *AJP* 2017; 174: 476-84.
51. Citrome L. Valbenazine for tardive dyskinesia: a systematic review of the efficacy and safety profile for this newly approved novel medication - What is the number needed to treat, number needed to harm and likelihood to be helped or harmed? *Int J Clin Pract* 2017; 71: e12964.
52. Giladi N. The mechanism of action of Botulinum toxin type A in focal dystonia is most probably through its dual effect on efferent (motor) and afferent pathways at the injected site. *J Neurol Sci* 1997; 152: 132-5.
53. Stahl SM. Comparing pharmacologic mechanism of action for the vesicular monoamine transporter 2 (VMAT2) inhibitors valbenazine and deutetrabenazine in treating tardive dyskinesia: does one have advantages over the other? *CNS Spectr* 2018; 23: 239-47.
54. Xu H, Qin H, Chen S, et al. Vitamin E in the treatment of tardive dyskinesia: a meta-analysis. *Int Clin Psychopharmacol* 2022; 37: 60-6.
55. Debrey SM, Goldsmith DR. Tardive dyskinesia: spotlight on current approaches to treatment. *FOC* 2021; 19: 14-23.
56. Caroff SN. A new era in the diagnosis and treatment of tardive dyskinesia. *CNS Spectr* 2023; 28: 401-15.
57. Hauser RA, Truong D. Tardive dyskinesia: out of the shadows. *J Neurol Sci* 2018; 389: 1-3.
58. Fernandez HH, Factor SA, Hauser RA, et al. Randomized controlled trial of deutetrabenazine for tardive dyskinesia: the ARM-TD study. *Neurology* 2017; 88: 2003-10.
59. Pouclet-Courtemanche H, Rouaud T, Thobois S, et al. Long-term efficacy and tolerability of bilateral pallidal stimulation to treat tardive dyskinesia. *Neurology* 2016; 86: 651-9.

Corresponding author:
Alessandro Cuomo
Policlinico S. Maria alle Scotte
Viale Mario Bracci 16
53100 Siena, Italy
E-mail: alessandrocuomo86@gmail.com

Appendix 1. Tardive dyskinesia: diagnosis and management: an expert opinion

AIMS examination procedure

Either before or after completing the Examination Procedure, observe the patient unobtrusively, at rest (e.g., in the waiting room). The chair to be used in this examination should be a hard, firm one without arms.

1. Ask the patient whether there is anything in his/her mouth (i.e., gum, candy, etc.) and if there is, remove it.
 2. Ask the patient about the current condition of his/her teeth. Do teeth bother the patient now?
 3. Ask the patient whether he/she notices any movements in his/her mouth, face, hands, or feet. If yes, ask the patient to describe them and to indicate to what extent they currently bother the patient or interfere with his/her activities.
 4. Have patient sit in chair with hands on knees, legs slightly apart, and feet flat on floor. (Look at entire body for movements while in this position).
 5. Ask patient to sit with hands hanging unsupported. If male, between legs; if female and wearing a dress, hanging over knees. (Observe hands or other body areas).
 6. Ask patient to open his/her mouth. (Observe the tongue at rest within the mouth). Do this twice.
 7. Ask patient to protrude his/her tongue. (Observe abnormalities of tongue movement). Do this twice.
 8. Ask patient to tap his/her thumb, with each finger as rapidly as possible for 10 to 15 seconds; first with right hand, then with left hand. (Observe facial and leg movements).
 9. Flex and extend patient's left and right arms (one at a time).
 10. Ask patient to stand up. (Observe the patient in profile. Observe all body areas again, hips included).
 11. Ask patient to extend both arms out in front, palms down. (Observe trunk, legs, and mouth).
 12. Have patient walk a few paces, turn, and walk back to the chair. (Observe hands and gait). Do this twice.
-

Administration time

- **Administration time:** the AIMS test generally takes about 10 to 15 minutes to complete. This includes both the interview and the physical examination of the patient.
-

Frequency of administration

- **Initial baseline measurement:** when first diagnosing or identifying symptoms of tardive dyskinesia, the AIMS should be administered to establish a baseline.
 - **Follow-up assessments:**
 - **Monthly:** for patients who are at a higher risk of developing TD (e.g., those on antipsychotic medications), it is recommended to conduct the AIMS monthly for the first year.
 - **Every 3 to 6 months:** after the first year, if the patient remains on the medication, the AIMS can be administered every 3 to 6 months. This helps to monitor for the emergence or progression of symptoms.
-

Clinical Judgment

- **More frequent monitoring:** in cases where there is a change in medication, an increase in dosage, or a change in the patient's clinical status, more frequent monitoring may be warranted.
- **Less frequent monitoring:** for patients who have been stable over a long period without any signs of TD, the frequency may be reduced based on clinical judgment.

Regular use of the AIMS allows healthcare providers to detect TD early and make necessary adjustments to the patient's treatment plan, potentially mitigating the severity of the disorder.

Abnormal Involuntary Movement Scale (AIMS)

Instructions: Complete the examination procedure before making ratings. Circle score for each item.

Patient Name:	Date:	None	Minimal, may be extreme normal	Mild	Moderate	Severe
Facial and Oral Movements						
1. Muscles of Facial Expression e.g., movements of forehead, eyebrows, periorbital area, cheeks; Include frowning, blinking, smiling, grimacing	0	1	2	3	4	
2. Lips and Perioral Area e.g., puckering, pouting, smacking	0	1	2	3	4	
3. Jaw e.g., biting, clenching, chewing, mouth opening, lateral movement	0	1	2	3	4	
4. Tongue Rate only increases in movement both in and out of mouth, NOT inability to sustain movement	0	1	2	3	4	
Extremity Movements						
5. Upper (arms, wrists, hands, fingers) Include choreic movements (i.e., rapid, objectively purposeless, irregular, spontaneous); athetoid movements (i.e., slow, irregular, complex, serpentine). DO NOT include tremor (i.e., repetitive, regular, rhythmic).	0	1	2	3	4	
6. Lower (legs, knees, ankles, toes) e.g., lateral knee movement, foot tapping, heel dropping, foot squirming, inversion and eversion of foot	0	1	2	3	4	
Trunk Movements						
7. Neck, shoulders, hips e.g., rocking, twisting, squirming, pelvic gyrations	0	1	2	3	4	
Global Judgments						
8. Severity of abnormal movements	0	1	2	3	4	
9. Incapacitation due to abnormal movements	0	1	2	3	4	
10. Patient's awareness of abnormal movements (rate only patient's report) 0 = not aware; 1 = aware, no distress; 2 = aware, mild distress; 3 = aware, moderate distress; 4 = aware, severe distress	0	1	2	3	4	
Dental Status						
11. Current problems with teeth and/or dentures?	No	Yes				

Appendix 2. Tardive dyskinesia: diagnosis and management: an expert opinion

Resume table.	
Step	Key points
Diagnosis	- Exposure to dopamine receptor blocking agents (DRBAs) for ≥3 months (≥1 month if age >60)
	- Presence of involuntary movements of tongue, face, lips, trunk, and extremities
	- Differential diagnosis to exclude other movement disorders
Assessment	- Use Abnormal Involuntary Movement Scale (AIMS) for screening and severity assessment
	- Consider Extrapyramidal Symptom Rating Scale (ESRS) for comprehensive evaluation
Treatment	- Regular screening: every 3-6 months for patients on first-generation antipsychotics, every 6-12 months for second-generation antipsychotics
	- Prevention: use lowest effective dose of antipsychotics
	- Consider discontinuation or dose reduction of offending agent if possible
	- Consider switching to clozapine for moderate evidence of symptom improvement - Consider use of a VMAT2 inhibitor - Explore non-pharmacological options (e.g., deep brain stimulation) for refractory cases